

vix distincto, apice emarginatis; *staminum filamentis omnibus appendiculatis*, *longiorum appendice subunilateralis ampla* in longitudine tota adnata *margine exteriore denticulata*, breviorum appendice oblonga superne libera integra; *siliculis ovato-suborbiculatis vel suborbiculatis*, apice subretusis, disco inflato valde convexis, margine complanatis, *glabris nitidulis*; *stylo brevi*; *seminibus anguste marginatis*. ①. Maio-Jun.

MAR. In glareosis et pascuis ad cacumen montis *Dj. Touchka* ad meridiem urbis *Maroc* [Ibr. 1875].

In montibus etiam excelsioribus Hispaniæ centralis et australis.

OBS. *A. psilocarpum* a cl. Boissier cum *A. calycino* comparatum ab eo plane distinctum glandulis valvariis minutis.

Var. β . *edentulum*. — *A. leiocarpum* Pomel *Nouv. mat.* 235 e descriptione. — Exs. Batt. in Soc. Dauph. *Alg.* 4460.

Staminum longiorum filamentis inferne dilatato-membranaceis margine exteriore integris.

In pascuis petrosis regionis Montanæ. — **Mont.** A¹: in declivitate australi cacuminis *Dj. Beni-Salah* ad 1600 m. (Batt.), *Dj. Zaccar* (sec. Pomel, l. c.).

OBS. E speciminiibus paucis suppetentibus, *A. leiocarpum* Pomel ad *A. psilocarpum*, cum quo habitu notisque plurimis exakte congruit, referendum et tanquam varietas tantum distinguendum filamentis omnibus staminum inferne plus minus dilatato-membranaceis margine exteriore integris.

6. A. macrocalyx Coss. et DR. ap. Coss. in *Bull. Soc. bot.* IV, 12, et *Illustr. Atl.* I, 60, t. 41. — Exs. War. *Alg.* 14.

أغورما Aguerma (Bertr.); *دريقة* *D'reiga* vel *D'ereiga*, diminutivum a *دُورج* *D'ourg* «nomen plantæ» (Lx).

Annum, stellato-pubescent; foliis oblongis vel oblongo-linearibus, inferne attenuatis; racemis demum saepius elongatis; *calyce subpersistente et post anthesim accrescente*, *sepalis crassiusculis conniventibus silicula matura paulo longioribus*; *petalis albis*, *calyce paulo longioribus*, linearibus ungue a limbo vix distincto, apice retusis vel integris; *staminum longiorum filamentis exappendiculatis*, *lateralium appendiculatis*; *siliculis majusculis*, *suborbiculatis*, apice vix retusis, disco inflato-convexis, margine complanatis, *glabris*; *stylo siliculae tertiam vel quartam partem subæquante*; *seminibus latiuscula submembranaceo-marginatis*. ①. Febr.-Maio.

In pascuis et depressis argilloso-arenosis vel glareosis, in alluviis et alveis, in Planitiebus excelsis australioribus, ubi saepius *A. scutigeri* socium, et margine contiguo regionis Saharensis. — **Plan.** A³: *Guelt-el-Settel* (Lx), *Djelfa* (Reb., Gesl.), *Oued Sedeur!*, inter *Rocher-de-Sel* et *Aïn-el-Ibel* (Lx), *Aïn-el-Ibel!*, *Sidi-Makhelouf!* (Vilm.), *Metlili!*. — O⁵: ex. gr.: *Stitten*, *Khodeur*, *Aïn-Tazina* (War.); *Bou-Guern*, *Dj. Antar*, *Mechera-el-Acheria*, *Chott El-Gharbi!* (War.), *Aïn-ben-Khelil!*, *Taous-*

sera!, etc. — **Sah.** A¹ : *Bou-Saada* (Reb., Soll.); *Aïn Milogh!*, *Aïn-Madhi!*, *Laghouat!* (Reb., Tess.), *Nili!* (Gesl., Vilm.), *Daïa El-Kebch!*; *Ghardaïa* (Vilm.). — O² : *Tadjerouna* (Par.), *Arba!*, inter *Chellala-Dahrania* et *Asla!*, *Bou-Semghoun* (War.), *Asla!* (War.), *Tyout!*, *Aïn-Sefra!*, *Aïn-Sefissifa!* (War.).

In Algeria hucusque tantum notum.

7. A. scutigerum DR. in *Atl. Alg.* t. 72, f. 4; Coss. et DR. ap. Coss. in *Bull. Soc. bot.* IV, 11. — *A. clypeatum* DR. olim in litt. — Exs. Bal. *Alg.* 536, Kral. 6, Choul. 205, War. 13, et in *Soc. Dauph.* 1087.

Annum, stellato-pubescent; foliis oblongis vel oblongo-linearibus, inferne attenuatis; racemis demum elongatis; *calyce* deciduo, *silicula* subdimidio breviore; *petalis albidis*, calyce paulo longioribus, oblongo-linearibus ungue a limbo vix distincto, apice emarginatis rarius integris; *staminum filamentis* omnibus appendiculatis; *siliculis* majusculis, suborbiculatis, apice retusis, disco inflato-convexis, margine complanatis, *glabris*; stylo siliculæ tertiam longitudinem subæquante; *seminibus latiuscule submembranaceo-marginatis*. ①. Mart.-Jun.

In pascuis et convallibus erosis præsertim calcareis vel argilloso-calcareis, in depressis hieme humidis, in alluviis et alveis, in Planitiebus excelsis unde nonnunquam hinc in valles regionis Montanæ et quidem Mediterraneæ, illinc in regionem Saharensem devehitur. — **Med.** O¹ : *Sidi-bel-Abbès* (Lefr.). — **Mont.** C² : *Dj. Mahmel!*, *Telet!*, *Haïdous!*, *Menah!*. — A² : *Dj. Afoul* prope *Aumale* (Char.); *Dj. Senalba!*. — O¹ : *Dj. Ksel!*. — **Plan.** C² : *Aïn Sedjera* (J. Reb.); *Aïn-Yagout!*, *Batna!* (Hén., Col., Kral., etc.), *Lambèse!*, *Ksour!*. — A⁵ : *Aumale*, in valle *Oued Chelif* prope *Boghar*, *Aïn-Oussera!*, *Guelt-el-Settel!*, *Djelfa!*, *Aïn-el-Ibel!*, etc. — O⁷. — **Sah.** A³ : *Bou-Saada* (Reb., Soll.), *Oued El-Anoug*, *Hassi-Dermel*, *Aïn-el-Ghorab*, *Aïn Melah* (Reb.); *Laghouat!*, *Tadjemout!*, *Aïn-Madhi!*. — O¹ : inter *Chellala-Gueblia* et *Arba-Tahtani!*. — **MAR.** Inter *Sebkha Tigri* et *Aïn-Chair* (Seign.), *Oued Bou-Roumad*, *Meharoug* (War.).

In Algeria et imperio Marocciano hucusque tantum obvium.

Subgen. III. TETRATRICHIA.

Tetratrichia J. Gay in herb. [1841]. — *Psilonema* C.-A. Mey. in *Led. Alt.* III, 50, et in *Led. Ross.* I, 137 emend.; *Endl. Gen.* n. 4873.

Petala lutescentia vel albida. Glandulæ hypogynæ ad basim utriusque staminis lateralis teretes subulatæ vel imo capillares. Staminum filimenta exappendiculata, rarius lateralium appendiculata. Silicula suborbiculata, valvis disco convexis, margine planis, loculis biovulatis. — Plantæ annuæ.

8. A. Granatense Boiss. et Reut. *Pug.* 9; Coss. *Pl. crit.* 146, et *Illustr. Atl.* I, 61, t. 42; Pomel *Nouv. mat.* 233; Willk. et Lange *Prodr. Hisp.* III, 833; Willk. *Illustr. Hisp.* I, t. 56, f. b. — *A. ptilotrichum* DR. in herb. olim. — *A. hispidum* Losc. var. *Granatense* in Willk. *Illustr. Hisp.* I, 85. — *A. Willkomii* de Rœm. ap. Willk.