




[Click for updates](#)

## Grana

Publication details, including instructions for authors and subscription information:  
<http://www.tandfonline.com/loi/sgra20>

# Pollen morphology of Ephedra (Gnetales) and its evolutionary implications

Kristina Bolinder<sup>a</sup>, Lena Norbäck Ivarsson<sup>b</sup>, Aelys M. Humphreys<sup>ac</sup>, Stefanie M. Ickert-Bond<sup>de</sup>, Fang Han<sup>f</sup>, Carina Hoorn<sup>g</sup> & Catarina Rydin<sup>a</sup>

<sup>a</sup> Department of Ecology, Environment and Plant Sciences, Stockholm University, Stockholm, Sweden

<sup>b</sup> School of Natural Sciences, Technology and Environmental Studies, Södertörn University, Huddinge, Sweden

<sup>c</sup> Department of Life Sciences, Imperial College London, Silwood Park Campus, Ascot, Berkshire, UK

<sup>d</sup> Department of Biology and Wildlife and Institute of Arctic Biology, University of Alaska Fairbanks, Fairbanks, AK, USA

<sup>e</sup> School of Life Sciences, Arizona State University, Tempe, AZ, USA

<sup>f</sup> State Key Laboratory of Biogeology and Environmental Geology, Faculty of Earth Science, China University of Geosciences, Wuhan, China

<sup>g</sup> Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, Amsterdam, The Netherlands

Published online: 18 Aug 2015.

To cite this article: Kristina Bolinder, Lena Norbäck Ivarsson, Aelys M. Humphreys, Stefanie M. Ickert-Bond, Fang Han, Carina Hoorn & Catarina Rydin (2015): Pollen morphology of Ephedra (Gnetales) and its evolutionary implications, Grana, DOI: [10.1080/00173134.2015.1066424](https://doi.org/10.1080/00173134.2015.1066424)

To link to this article: <http://dx.doi.org/10.1080/00173134.2015.1066424>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

## Pollen morphology of *Ephedra* (Gnetales) and its evolutionary implications

KRISTINA BOLINDER<sup>1</sup>, LENA NORBÄCK IVARSSON<sup>2</sup>, AELYS M. HUMPHREYS<sup>1,3</sup>,  
STEFANIE M. ICKERT-BOND<sup>4,5</sup>, FANG HAN<sup>6</sup>, CARINA HOORN<sup>7</sup> & CATARINA RYDIN<sup>1</sup>

<sup>1</sup>Department of Ecology, Environment and Plant Sciences, Stockholm University, Stockholm, Sweden, <sup>2</sup>School of Natural Sciences, Technology and Environmental Studies, Södertörn University, Huddinge, Sweden, <sup>3</sup>Department of Life Sciences, Imperial College London, Silwood Park Campus, Ascot, Berkshire, UK, <sup>4</sup>Department of Biology and Wildlife and Institute of Arctic Biology, University of Alaska Fairbanks, Fairbanks, AK, USA, <sup>5</sup>School of Life Sciences, Arizona State University, Tempe, AZ, USA, <sup>6</sup>State Key Laboratory of Biogeology and Environmental Geology, Faculty of Earth Science, China University of Geosciences, Wuhan, China, <sup>7</sup>Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, Amsterdam, The Netherlands

### Abstract

The *Ephedra* lineage can be traced at least to the Early Cretaceous. Its characteristically polyplicate pollen is well-represented in the fossil record and is frequently used as an indicator of paleoclimate. However, despite previous efforts, knowledge about variation and evolution of ephedroid pollen traits is poor. Here, we document pollen morphology of nearly all extant species of *Ephedra*, using a combination of scanning electron microscopy (SEM) and light microscopy (LM), and reconstruct ancestral states of key pollen traits. Our results indicate that the ancestral *Ephedra* pollen type has numerous plicae interspaced by unbranched pseudosulci, while the derived pollen type has branched pseudosulci and (generally) fewer plicae. The derived type is inferred to have evolved independently twice, once along the North American stem branch and once along the Asian stem branch. Pollen of the ancestral type is common in Mesozoic fossil records, especially from the Early Cretaceous, but it is less commonly reported from the Cenozoic. The earliest documentation of the derived pollen type is from the latest Cretaceous, after which it increases strongly in abundance during the Paleogene. The results of the present study have implications for the age of crown group *Ephedra* as well as for understanding evolution of pollination syndromes in the genus.

**Keywords:** character evolution, light microscopy, phylogeny, polyplicate, pseudosulci, scanning electron microscopy, Welwitschia

Pollen plays an important role in the lifecycle of all seed plants. Consequently, pollen characters have shown to be informative in studies of plant evolution and for resolving phylogenies (e.g. Doyle & Le Thomas 1994; Doyle & Endress 2000; Sauquet & Le Thomas 2003), for calibrating molecular dating analyses (Thornhill et al. 2012), as well as for studying plant reproductive biology (Ferguson & Skvarla 1982; Grayum 1986; Osborn et al. 1991; Bolinder et al. 2015). In addition, fossil pollen data are also frequently used for reconstructing past vegetation

types and for inferring paleoclimates (Hoorn et al. 2012). Ephedroid pollen (i.e. pollen inferred to have been produced by *Ephedra* (Gnetales) or *Ephedra*-like extinct plants) is characteristically polyplicate, well known from the fossil record, and considered a good indicator of a very dry paleoclimate (Li et al. 2005; Hoorn et al. 2012).

The earliest reported pollen of probable ephedroid affinity dates to the Permian (Wilson 1962; Wang 2004). By the Early Cretaceous, ephedroid pollen had rapidly increased in abundance and distribution

Correspondence: Kristina Bolinder, Department of Ecology, Environment and Plant Sciences, Stockholm University, SE-106 91 Stockholm, Sweden.  
E-mail: [Kristina.Bolinder@su.se](mailto:Kristina.Bolinder@su.se)

(Received 7 April 2015; accepted 11 June 2015)

© 2015 Collegium Palynologicum Scandinavicum




Figure 1. Schematic drawing of the polarity and different types of pseudosulci branching in *Ephedra* pollen. The long equatorial axis is equal to the longest axis and the polar axis is equal to one of the shortest axes. **A.** Pollen with unbranched pseudosulci; the ancestral type. **B.** Pollen with pseudosulci with first-order branching; the derived type. **C.** Pollen with pseudosulci with first- and second-order branching; the derived type.

at low palaeo-latitudes (Crane & Lidgard 1989), and a relatively large diversity of ephedroid megafossils is described from this time (e.g. Krassilov 1986; Yang et al. 2005; Rydin et al. 2006b; Wang & Zheng 2010). Towards the latter part of the Cretaceous, however, the abundance of ephedroid pollen appears to have decreased substantially (Crane & Lidgard 1989). The fossil record after the K-Pg boundary is perhaps less well studied, but there are currently no reported megafossils of ephedroid affinity after the Early Cretaceous. Current knowledge of the Cenozoic diversity of *Ephedra* (before the present) is thus based on the palaeo-palynological record, which is extensive and present also at higher palaeo-latitudes than are Cretaceous records (Akkiraz et al. 2008; Wodehouse 1933; Cookson 1956; Gray 1960; Ghosh et al. 1963; Nagy 1963; Shaw 1998; Hoorn

et al. 2012). Today, *Ephedra* comprises about 50 species with remarkably low genetic (Ickert-Bond & Wojciechowski 2004; Rydin & Korall 2009) and morphological (Rydin et al. 2010) diversity, occupying arid regions of the world (Kubitzki 1990). Molecular analyses imply that the clade of extant species dates to the earliest Oligocene (Ickert-Bond et al. 2009; Rydin et al. 2010), and it has been suggested that most of this diversity might be the result of radiation caused by a shift from insect-pollination to wind-pollination early in the evolution of the crown group (Bolinder et al. 2014).

*Ephedra* pollen is large, between 34 and 81  $\mu\text{m}$  in its longest (equatorial) diameter (Steeves & Barghoorn 1959), and, as in remaining Gnetales, the pollen wall consists of a homogenous tectum, a granular infratectum of varying density, and a thin foot layer adnate to a distinct lamellar endexine (Gullvåg 1966; Van Campo & Lugardon 1973; Hesse 1984; Zavada 1984; Kurmann 1992; Rowley 1995; El-Ghazaly & Rowley 1997; Osborn 2000; Tekleva & Krassilov 2009; Bolinder et al. 2015). Based on developmental studies, Huynh (1975) and El-Ghazaly et al. (1998) concluded that the longest axis in *Ephedra* pollen is equatorial and the polar axis is equal to one of the shortest axes (Huynh 1975; El-Ghazaly et al. 1998) (Figure 1). Although *Ephedra* pollen is typically described as inaperturate (Erdtman 1952; Huynh 1975; Kurmann & Zavada 1994; El-Ghazaly et al. 1998; Ickert-Bond et al. 2003; Doores et al. 2007), some authors have interpreted *Ephedra* pollen as polyaperturate (Steeves & Barghoorn 1959; Bharadwaj 1963), referring to the furrows that run between the plicae parallel to the long equatorial axis. In these furrows, which have been called hyaline lines (Woodhouse 1935; Steeves & Barghoorn 1959; Kedves 1987; Kurmann & Zavada 1994; El-Ghazaly et al. 1998), pseudosulci (Huynh 1975; Bolinder et al. 2015) and colpi (Steeves & Barghoorn 1959; Zhang & Xi 1983), the exine is much thinner than over the ridges and neither the tectum nor the infratectum is present (Osborn 2000; Tekleva & Krassilov 2009; Bolinder et al. 2015). When the pollen germinates, the exine splits open in two of these furrows and detaches from the intine (Land 1907; Mehra 1938; El-Ghazaly et al. 1998), and, based on the polarity described by Huynh (1975) and El-Ghazaly et al. (1998), we will hereafter refer to the furrows as pseudosulci (Figure 1) (following Huynh 1975; Bolinder et al. 2015).

Woodhouse (1935) classified *Ephedra* pollen into two types based on the number of ridges and the appearance of the 'hyaline line' in the grooves (i.e. the pseudosulcus). Later, Steeves and Barghoorn (1959) divided *Ephedra* pollen into four

groups (Type A–D) based on the number and appearance of the ridges as well as the presence or absence of ‘colpi’. Their type A has an average of five to nine plicae, which are triangular in the transverse section and interspaced by narrowly serpentine, sometimes laterally branched colpi. Pollen of type B has indistinct colpi and an average of 10 to 13 plicae, and type C is similar to B but with higher plicae as seen in the transverse section. Type D shows numerous plicae, up to 20, which are wide and rounded in the transverse section and not interspaced by colpi (Steeves & Barghoorn 1959). Zhang and Xi (1983) merged types B and C of Steeves and Barghoorn (1959), thus recognising three pollen types. In line with Woodhouse (1935), Kedves (1987) and Freitag and Maier-Stolte (1994) recognised only two pollen forms, based on the presence or absence of a ‘hyaline line’.

Extensive intraspecific variation and dimorphism in pollen morphology have been reported (El-Ghazaly & Rowley 1997; Ickert-Bond et al. 2003; Doores et al. 2007). It has also been shown recently that the morphology and ultrastructure differ between *Ephedra* pollen of anemophilous and entomophilous species and that these differences influence the aerodynamic properties of the pollen grains (Bolin-der et al. 2015). Although pollen of living species of *Ephedra* has been studied previously, few studies have aimed to assess pollen morphology across the entire genus. More importantly, *Ephedra* pollen morphology has never been studied in an evolutionary context. We use a combination of scanning electron microscopy (SEM) and light microscopy (LM) and a much larger sample than previously utilised to study variation and evolution of pollen morphology in extant *Ephedra* and compare the results with available information from ephedroid fossil pollen. We also investigate whether it is possible to assign extant and fossil pollen to specific subclades or species of extant *Ephedra*.

## Material and methods

### *Taxon sampling*

We selected 45 species for the present study, spanning the phylogenetic and geographical diversity of the genus and representing about 85% of the species. Pollen from two to five specimens of each species was studied, except for a few species (*Ephedra alata*, *E. aspera*, *E. boelckei*, *E. compacta* and *E. trifurcata*), where limited access to material prevented study of more than one specimen per species. The specimens studied by Steeves and Barghoorn (1959) are currently deposited at the herbaria (A) and (GH) and all specimens still available were included in the present

study. For a full list of herbarium accessions, see the ‘Specimens investigated’ section.

### *Sample preparation*

Both the size and morphology of Gnetales pollen are affected by conventional preparation methods (i.e. acetolysis and staining; Kedves 1987). Therefore, we tested if treatment in alcohol can have the same effect, and if pollen extracted directly from herbarium sheets, without further treatment, is a suitable way of studying natural variation in *Ephedra* pollen. For this purpose, anthetic microsporangiate structures of *E. viridis* were harvested from living plants housed in the glasshouses at Stockholm University. Pollen grains obtained from these plants were treated in five different ways before study: (a) no treatment and examination within one hour of collection ( $n = 30$ ), (b) air-dried in an envelope for a week to approximate herbarium-dried material ( $n = 30$ ), (c) placed in 70% ethanol ( $n = 30$ ), (d) placed in 70% ethanol followed by dehydration with a conventional ethanol series ( $n = 30$ ), and (e) air-dried for a week to approximate herbarium-dried material, washed with phosphate-buffered saline (PBS) and dehydrated in an ethanol series prior to investigation ( $n = 30$ ). Following this preparation, the size and morphology of pollen grains were compared among treatments using an analysis of variance (ANOVA) and Tukey’s Honestly Significant Differences (HSD) test in R version 3.1. (R Development Core Team 2014). Having established the best way to study natural variation in pollen morphology, anthetic structures were obtained from herbarium material and pollen grains were studied using SEM and LM without any treatment or preparation prior to the investigations.

### *Scanning electron microscopy and light microscopy*

A minimum of 20 pollen grains per species were obtained from herbarium material and mounted on aluminium stubs using double-sided tape, sputter-coated with gold (40 s at 10 mA) and studied under SEM. Abnormal and seemingly aborted pollen was carefully avoided. In addition, one representative specimen of each species was selected for comparative studies using LM. For this purpose, 15 pollen grains of each species were mounted in glycerine and studied under LM with a 40× objective.

The lengths of polar and longest equatorial axes were measured during the SEM studies, and a shape estimate (polar axis/equatorial axis, P/E-ratio) was obtained. The number of plicae was counted on the visible side of the grain and multiplied by two to obtain the total number of plicae. Presence or

absence of first- and second-order branches on the pseudosulci (Figure 1) was scored for each pollen grain.

#### *Statistical comparison of pollen characters*

Appropriate sample size for each species was controlled according to Van Emden (2008) at 95% power to limit the risk of Type I errors. To determine the best way to analyse the pollen data, the phylogenetic signal in each of the variables was first estimated using the branch length transformation parameter Pagel's lambda,  $\lambda$  (Pagel 1999a), a robust index with low Type I error rates (Freckleton et al. 2002). This allowed assessment of the extent to which interspecific differences in pollen traits are correlated with phylogenetic relatedness. The parameter  $\lambda$  may vary between 1 (if trait variance is perfectly correlated with phylogenetic distance, equivalent to a Brownian Motion model (BM; Schluter et al. 1997) and 0 (there is no relationship between trait variance and phylogenetic distance). Alternatively,  $\lambda$  may assume an intermediate value if there is some degree of phylogenetic dependence in the trait ( $0 < \lambda < 1$ ). The value of  $\lambda$  was determined by comparing the likelihood fit of three different models to each pollen variable (1:  $\lambda = 1$ , 2:  $\lambda = 0$  and 3:  $\lambda$  is estimated during model fitting) as implemented in the R package *motmot* (Thomas & Freckleton 2012). Model fit was assessed using the Akaike Information Criterion (AIC; Akaike 1974). The pollen variables were absolute size, as assessed by the long equatorial diameter, pollen shape, as gauged by the P/E ratio, and the number of plicae. Phylogenetic information was obtained from Rydin and Korall (2009).

In characters for which no correlation with the phylogeny was established (i.e. the best model is when  $\lambda = 0$ ), variation was compared within and among species and clades (as defined in Figures 13 and 14 later) using a conventional ANOVA, Tukey's HSD and model selection using AIC in R (R Development Core Team 2014). For characters, where variance is correlated with the phylogeny (i.e.  $\lambda$  is significantly different from 0), variation was not compared any further.

#### *Parsimony ancestral state reconstruction of pollen characters*

Parsimony reconstruction of ancestral states for each pollen character was conducted based on to date the most well-sampled phylogeny of *Ephedra* (Rydin & Korall 2009), using the Trace Character History command in Mesquite version 2.75 (Maddison & Maddison 2011). These analyses were performed in

order to assess the relative amount of evolutionary information in pollen characters. Parsimony was considered appropriate for this because, although it is appropriate for detailed analyses only when transition rates are low (Harvey & Pagel 1991; Pagel 1999b; Pirie et al. 2012), it will still allow determination of which pollen characters display a more conserved phylogenetic pattern. Ancestral states were estimated for the presence or absence of side branches on the pseudosulci using two alternative codings: (a) a binary state option: 0, absence of side branches on the pseudosulci (Figure 1A); or 1, presence of side branches on the pseudosulci (Figure 1B, C), and (b) a multistate approach: 0, complete absence of side branches on the pseudosulci (Figure 1A); 1, presence of first-order side branches (Figure 1B); and 2, presence of first- and second-order side branches on pseudosulci (Figure 1C). To compare observed estimates of gains and losses to those expected by chance, the terminals were shuffled 999 times using the Reshuffle Character command and the character history of each reshuffled character was traced. This allowed a distribution of parsimony steps needed for random characters to be compared with the observed number of steps.

For the continuous characters (number of plicae, length of the polar axis and the P/E ratio), the correlation between minimum, maximum and mean values was tested for, using simple regression with a linear model in R (R Development Core Team 2014). Minimum, maximum and mean values are strongly correlated (number of plicae:  $r^2 = 0.96$ ,  $p \ll 0.05$ ; length of the long equatorial axis:  $r^2 = 0.87$ ,  $p \ll 0.05$ ; P/E ratio:  $r^2 = 0.77$ ,  $p \ll 0.05$ ); therefore, mean values were used to reconstruct ancestral states using parsimony as earlier.

## **Results**

### *Sample preparation*

*Size (length of the long equatorial axis).* — There is a significant difference in the length of the long equatorial axis among pollen grains treated in different ways prior to investigation ( $F_{145, 4} = 65.4$   $p \ll 0.05$ ). However, there is no significant difference between freshly collected and air-dried pollen grains (Tukey's HSD;  $p = 0.16$ ). Pollen obtained from herbarium material therefore captures the natural variation in pollen size in *Ephedra* (Figure 2A). There is a significant difference between pollen grains treated with ethanol in various ways compared to fresh and air-dried (Tukey's HSD; fresh versus ethanol  $p \ll 0.05$ ; fresh versus air-dried + dehydration series  $p \ll 0.05$ ; fresh versus ethanol + dehydration series  $p \ll 0.05$ ;

air-dried versus ethanol + dehydration series  $p \ll 0.05$ ; fresh versus air-dried + dehydration series  $p \ll 0.05$ ; air-dried versus air-dried + dehydration series  $p \ll 0.05$ ).

**Shape (P/E ratio).** — There is a significant difference in the P/E ratio among pollen grains treated in different ways ( $F_{145, 4} = 27.6$   $p \ll 0.05$ ). However, there is no significant difference between fresh and air-dried pollen grains (Tukey's HSD;  $p = 0.46$ ; Figure 2B). Further, there is no difference between pollen grains subjected to the different ethanol treatments (Tukey's HSD; ethanol + dehydration series versus ethanol  $p = 0.06$ ; air-dried + dehydration series versus ethanol  $p = 0.75$ ; ethanol + dehydration series versus air-dried + dehydration series  $p = 0.55$ ). A significant difference in P/E ratio between pollen grains treated with ethanol in various ways compared to fresh and air-dried pollen grains (Tukey's HSD; fresh versus ethanol  $p \ll 0.05$ ; air-dried + dehydration series versus fresh  $p \ll 0.05$ ; fresh versus ethanol + dehydration series  $p \ll 0.05$ ; air-dried versus ethanol + dehydration series  $p \ll 0.05$ ; fresh versus air-dried + dehydration series  $p \ll 0.05$ ; air-dried versus air-dried + dehydration series  $p \ll 0.05$ ).

#### Pollen morphology

**Mediterranean species (Figures 3, 7; Table I).** — Pollen of Mediterranean species has 10 to 22 plicae extending parallel to the long equatorial axis and fusing at the tips. The plicae are psilate, wide and rounded in transverse section. Between adjacent plicae, there is a distinct or indistinct, unbranched pseudosulcus.

**North American species (Figures 4, 8; Table I).** — Pollen of *Ephedra californica*, *E. trifurca*, *E. pedunculata* and *E. torreyana* has 10 to 22 plicae that extend parallel to the long equatorial axis and fuse at the tips. The plicae are psilate, wide and rounded in transverse section. Between adjacent plicae there is a distinct or indistinct, unbranched pseudosulcus. Pollen of *E. antisiphilitica* and *E. compacta* has distinct pseudosulci that occasionally have first-order branches. Pollen of *E. aspera*, *E. fasciculata* and *E. funerea* has 5 to 18 narrow plicae that are psilate and triangular in transverse section and the pseudosulci have first-order branching. Pollen of *E. coryi*, *E. cutleri*, *E. nevadensis* and *E. viridis* has 4 to 12 plicae and the pseudosulci are always branched, often with both first- and second-order branches.

**South American species (Figures 5, 9; Table I).** — Pollen of South American species has 10 to 22 plicae that extend parallel to the long equatorial axis and fuse




Figure 2. Effect of preparation method on two pollen traits. **A.** Length of the long equatorial axis (in micrometres). There are significant differences between pollen grains treated with ethanol in various ways and untreated pollen. There is no significant difference between fresh and air-dried pollen grains. **B.** P/E ratio. There are significant differences between pollen grains treated with ethanol and untreated pollen. There is no significant difference between fresh and air-dried pollen grains.

at the tips. The plicae are psilate, wide and rounded in transverse section. Between adjacent plicae, there is a distinct or indistinct, unbranched pseudosulcus.

**Asian species (Figures 6, 10; Table I).** — Pollen of Asian species generally has 4 to 12 plicae that extend parallel to the long equatorial axis and fuse at the tips. The plicae are psilate, narrow and triangular in transverse section. Between adjacent plicae there is a distinct, branched pseudosulcus that often shows first- and second-order branches. Pollen of *Ephedra likangensis*, *E. lomatolepis*, *E. minuta* and *E. saxatilis*




Figure 3. Pollen of *Ephedra*, Mediterranean species, scanning electron micrographs. **A.** *E. foeminea*, R Pampino & R Pichi-Sermolli 139 (L). **B.** *E. alata* AA Anderberg 480 (S). **C.** *E. altissima* H Freitag 35035 (KAS). **D.** *E. aphylla* G Samuelsson 2696 (S). **E.** *E. ciliata* Handel-Mazetti 973 (WU). **F.** *E. foliata* Hedberg & Hedberg 92019A (UPS). **G.** *E. milleri* Miller 7667A (E). Scale bars – 10  $\mu\text{m}$ .

has 10 to 20 plicae and branched pseudosulci (first-order branches only), deviating from the general pattern. The same holds for pollen of *E. sarcocarpa*, *E. strobilacea* and *E. transitoria*, which has 10 to 22 plicae and pseudosulci that are never branched.

#### Statistical comparison of pollen characters among species

In *Ephedra*, variance in the number of plicae is correlated with phylogenetic distance, i.e. the lambda model showed the best fit ( $\lambda = 0.61$ ; 95% confidence interval: 0.23–0.86;  $\Delta\text{AIC} = 30.4$  compared to BM and  $\Delta\text{AIC} = 11.7$  compared to the model with  $\lambda = 0$ ). It is clear that number of plicae varies considerably among species (Figure 12A) as well as among the different clades and pollen types (Figures 11A, B).

Variation among species in both pollen size (as gauged by the long equatorial diameter) and pollen shape (estimated by the P/E ratio) is independent of phylogeny. For both variables, this model ( $\lambda = 0$ ) was much better than BM ( $\Delta\text{AIC} = 24.4$  [pollen size],  $\Delta\text{AIC} = 33.4$  [pollen shape]) and indistinguishable from the lambda model ( $\Delta\text{AIC} = 0.0$  [pollen size],  $\Delta\text{AIC} = 1.21$  [pollen shape]). Consequently, we used a conventional one-way ANOVA of independent groups, Tukey's HSD and model selection using AIC to compare pollen size and pollen shape within and among species and clades.

The size varies significantly among the different clades  $F_{3,4018} = 32.5$ ,  $p \ll 0.05$ ; only the North American clade (hereafter referred to as defined in Figures 13 and 14, excluding *Ephedra pedunculata*) do not differ significantly from Mediterranean




Figure 4. Pollen of *Ephedra*, North American species, scanning electron micrographs. **A.** *E. pedunculata* RM Stewart 2265 (GH). **B.** *E. compacta* DS Corell & IM Johnston 20233 (NY). **C.** *E. viridis* LS Rose 58080 (S). **D.** *E. trifurca* CV Hartman 642 (GH). **E.** *E. californica* A Carter 3667 (L). **F.** *E. fasciculata* JH Lehr 2309 (NY). **G.** *E. aspera* BA Stein 31 (RSA). **H.** *E. nevadensis* P Raven 14251 (WU). **I.** *E. cutleri* CT Mason Jr. 2192 (TEX). **J.** *E. torreyana* Spellenberg 10204 (TEX). **K.** *E. funerea* CL Hitchcock 329 (GH). **L.** *E. antisiphilitica* Henderson 62-02a (BR). **M.** *E. coryi* DS Corell 32785 (S). Scale bars – 10  $\mu\text{m}$ .

species ( $p = 0.41$ ) (Figure 11B). Pollen size also differs significantly among species  $F_{3,3977} = 46.46$ ,  $p \ll 0.05$ , as well as within some of the species (Figure 12B). However, the variation in size among species is significantly larger than among clades or within species ( $\Delta\text{AIC} = 1489$  [among clades],  $\Delta\text{AIC} = 0$  [among species],  $\Delta\text{AIC} = 1572$  [within species]). Also, pollen shape differs significantly among the different clades  $F_{3,4018} = 48.43$ ,  $p \ll$

0.05; but North American species do not differ significantly from South American species ( $p = 0.41$ ) or Mediterranean species ( $p = 0.09$ ) (Figure 11E). Pollen shape varies significantly among species  $F_{44,3977} = 26.06$ ,  $p \ll 0.05$  (Figure 12C), but the variation in shape is greater among species than among clades or within species ( $\Delta\text{AIC} = 794$  [among clades],  $\Delta\text{AIC} = 0$  [among species],  $\Delta\text{AIC} = 930$  [within species]).




Figure 5. Pollen of *Ephedra*, South American clade, scanning electron micrographs. **A.** *E. americana* Gerth s.n. (L). **B.** *E. breana* KH & W Rechinger 63547 (W). **C.** *E. chilensis* C. Skottsberg 987 (F). **D.** *E. frustillata* MP Moreno 236 (NY). **E.** *E. multiflora* J Chiapella 2344 & E Vitek 09–0359 (W). **F.** *E. boelckeii* Maas et al. 8184 (GB). **G.** *E. ochreatea* RH Fortunato 5413 (NY). **H.** *E. triandra* J Chiapella 2505 & E Vitek 09–0520 (W). **I.** *E. trifurcata* O Zöllner 7928 (L). **J.** *E. tweediana* JH Hunziker 1648 (S). Scale bars – 10  $\mu\text{m}$ .

#### *Ancestral state reconstruction of discrete pollen characters*

*Presence of side branches on pseudosulci.* — There are five parsimony steps in the observed data compared to 7–20 for randomised data. Thus, the observed estimate does not overlap with the distribution of estimates for the random traits. Side branches are inferred to have evolved twice, once along the stem branch of the Asian clade and once along the stem branch of the North American species except *Ephedra pedunculata*. Side branches

have subsequently been lost once among Asian species and twice among North American species (Figure 13A) and some species show a reversal to the ancestral state of having no side branches. Dividing the character further, and discriminating between having first-order branching only and first- and second-order branching of pseudosulci, 12 steps are inferred for the observed data compared to 12–20 for the randomised data. Thus, the observed estimate overlaps with the distribution of estimates for the random traits at the




Figure 6. Pollen of *Ephedra*, Asian clade, scanning electron micrographs. **A.** *E. likiangensis* G Forrest 5564 (BM). **B.** *E. minuta* B Dikoré 8457 (MSB). **C.** *E. equisetina* QR Wu (MO). **D.** *E. gerardiana* Walter Koelz 5310 (S). **E.** *E. monosperma* H Freitag 33068 (KAS). **F.** *E. pachyclada* KH & F. Rechinger 3676 (W). **G.** *E. saxatilis* Parkinson 7077 (S). **H.** *E. intermedia* Lindberg 117–1947 (W). **I.** *E. lomatolepis* II Rusanovick & LA Krai 56 (NY). **J.** *E. regeliana* H Hastman 26 (MSB). **K.** *E. sinica* Y Yang 99531 (PE). **L.** *E. distachya* J Prudhomme 89 (WU). **M.** *E. sarcocarpa* KH Rechinger 46054b (W). **N.** *E. strobilacea* KH & F Rechinger 2703 (S). **O.** *E. transitoria* Doppelbaur 190 (M). Scale bars – 10  $\mu\text{m}$ .

99th percentile. Second-order branching is inferred to have evolved once in the Asian clade and to have been lost twice. In addition, side branches have been entirely lost in a small clade, comprising *E. strobilacea*, *E. sarcocarpa* and *E. transitoria*. In the clade comprising the North American species except *E. pedunculata*, second-order branching is inferred to have originated at least once and to have been lost again at least once. In addition, and as in the Asian clade, reversals back to unbranched pseudosulci have occurred several times among the North American species (Figure 13B).

#### *Ancestral state reconstruction of continuous pollen characters*

*Mean number of plicae.* — The ancestral state in *Ephedra* is numerous plicae (14.8–16.14). During the course of evolution, there has been a general trend towards fewer plicae in all clades, but a few species are inferred to have evolved an even greater number of plicae (up to about 20; Figure 14A).

*Mean length of the long equatorial axis (size).* — There is ample size variation within and among species (Figure 12B) and most species have a mean equatorial




Figure 7. Pollen of *Ephedra*, Mediterranean species, light micrographs. **A.** *E. foeminea* R Pampino & R Pichi-Sermolli 139 (L). **B.** *E. alata* AA Anderberg 480 (S). **C.** *E. altissima* H Freitag 35035 (KAS). **D.** *E. aphylla* G Samuelsson 2696 (S). **E.** *E. ciliata* Handel-Mazetti 973 (WU). **F.** *E. foliata* Hedberg & Hedberg 92019A (UPS). **G.** *E. milleri* Miller 7667A (E). Scale bars – 10  $\mu\text{m}$ .

diameter ranging between 32.8 and 49.8  $\mu\text{m}$ , although a few unrelated species stand out as having smaller (*Ephedra lomatolepis*, *E. boelckeii* and *E. funerea*) or larger (*E. alata*, *E. sarcocarpa*, *E. viridis* and *E. nevadensis*) pollen grains.

*Mean P/E ratio (shape).* — The ancestral state in *Ephedra* is a P/E ratio of 0.42 to 0.44, a state shared by most Mediterranean, South American and North American species. Among Asian species, a somewhat smaller P/E ratio (0.39–0.41) is more common (Figure 14C).

## Discussion

### Pollen morphology

*Pseudosulci.* — The appearance of the pseudosulci is the most important pollen morphological difference among species of *Ephedra*, and perhaps also the most important character from an ecological and evolutionary perspective. The ancestral pollen type in *Ephedra* lacks side branches of the pseudosulci, whereas pollen with branched pseudosulci represents a derived pollen type. The only *Ephedra* species known to be insect-pollinated (*E. foeminea*, Bolinder et al. 2014, and perhaps also *E. aphylla*, Bino et al.




Figure 8. Pollen of *Ephedra*, North American species, light micrographs. **A.** *E. pedunculata* RM Stewart 2265 (GH). **B.** *E. compacta* DS Corell & I. M. Johnston 20233 (NY). **C.** *E. viridis* LS Rose 58080 (S). **D.** *E. trifurca* CV Hartman 642 (GH). **E.** *E. californica* A Carter 3667 (L). **F.** *E. fasciculata* JH Lehar 2309 (NY). **G.** *E. aspera* BA Stein 31 (RSA). **H.** *E. nevadensis* Raven 14251 (WU). **I.** *E. cutleri* CT Mason jr. 2192 (TEX). **J.** *E. torreyana* Spellenberg 10204 (TEX). **K.** *E. funerea* CL Hitchcock 329 (GH). **L.** *E. antisiphilitica* Henderson 62-02a (BR). **M.** *E. coryi* DS Corell 32785 (S). Scale bars – 10  $\mu$ m.

1984) show the ancestral pollen type (Bolinder et al. 2014, 2015; Rydin & Bolinder 2015) and has a denser ultrastructure and therefore a reduced flight capacity compared to the derived pollen type (Bolinder et al. 2015). However, the correlation between

pollination syndrome and pollen morphology is ambiguous, because pollen of some putative anemophilous species (i.e. *E. trifurca*; Niklas et al. 1986; Niklas & Kerchner 1986; Buchmann et al. 1989; Niklas 2015) is also of the ancestral pollen type.




Figure 9. Pollen of *Ephedra*, South American clade, light micrographs. **A.** *E. americana* Gerth s.n. (L). **B.** *E. breana* KH & W Rechinger 63547 (W). **C.** *E. chilensis* C. Skottsberg 987 (F). **D.** *E. frustillata* MP Moreno 236 (NY). **E.** *E. multiflora* J Chiapella 2344 & E Vitek 09–0359 (W). **F.** *E. boelckeii* Maas et al. 8184 (GB). **G.** *E. ochreata* RH Fortunato 5413 (NY). **H.** *E. triandra* J Chiapella 2505 & E Vitek 09–0520 (W). **I.** *E. trifurcata* O Zöllner 7928 (L). **J.** *E. tweediana* JH Hunziker 1648 (S). Scale bars – 10  $\mu$ m.

Branched pseudosulci (the derived pollen type) appear to have evolved twice independently, once along the branch leading to the Asian clade and once along the branch leading to the North American clade (excluding *Ephedra pedunculata*; Figure 13). This indicates that there might, in fact, be two separate kinds of derived pollen types in *Ephedra*; however, we find it impossible to distinguish the derived pollen type of Asian species from that of North American species, suggesting, instead, convergent evolution of identical pollen types. South American species, as well as the North American

*E. pedunculata*, appear to have retained the ancestral pollen type, and their pollen cannot be distinguished from that of the Mediterranean species or from a few Asian and other North American species. It is thus not possible to identify species based solely on pollen morphology; not even assignment to a particular subclade of *Ephedra* is possible. For example, although first-order side branches on the pseudosulci are nearly universally present among North American and Asian species, the feature has been lost once in the Asian clade and at least twice in the North American clade (Figure 13A). Second-order





Figure 10. Pollen of *Ephedra*, Asian clade, light micrographs. **A.** *E. likiangensis* G Forrest 5564 (BM.) **B.** *E. minuta* B Dikoré 8457 (MSB). **C.** *E. equisetina* QR Wu (MO). **D.** *E. gerardiana* Walter Koelz 5310 (S). **E.** *E. monosperma* H Freitag 33068 (KAS). **F.** *E. pachyclada* KH. & F Rechinger 3676 (W). **G.** *E. saxatilis* Parkinson 7077 (S). **H.** *E. intermedia* Lindberg 117–1947 (W). **I.** *E. lomatolepis* II Rusanovick & LA Krai 56 (NY). **J.** *E. regeliana* H Hastman 26 (MSB). **K.** *E. sinica* Y Yang 99531 (PE). **L.** *E. distachya* J Prudhomme 89 (WU). **M.** *E. sarcocarpa* KH Rechinger 46054b (W). **N.** *E. strobilacea* KH & F Rechinger 2703 (S). **O.** *E. transitoria* Doppelbaur 190 (M). Scale bars – 10  $\mu$ m.

branching of the pseudosulci is not a good characteristic to assign pollen to clades or identify monophyletic groups in *Ephedra* either: It is inferred to have evolved at least twice, once in the Asian clade and once among the North American species, and to have later been lost repeatedly (Figure 13B). It should be noted that the presented analyses do not accommodate for phylogenetic uncertainty, however, we find it impossible to distinguish among pollen types with second-order branching, regardless of whether the pollen comes from closely related species or from species of different clades and

geographical regions, and the character cannot be used as a diagnostic feature.

The function of branched pseudosulci and/or second-order branches is not fully understood. In general, pollen morphology is not only related to pollination mode but also to the degree of dehydration at dispersal (Franchi et al. 2002). Pollen of species that release partly hydrated pollen has a thick pollen exine, more rapid germination of the male gametophyte and longer pollen viability (Franchi et al. 2002). In *Ephedra*, pollen with unbranched pseudosulci (the ancestral type) has a thick exine and



germinates faster than pollen with branched pseudosulci (the derived types; Bolinder et al. 2015 and KB personal observation, June 2012). Although this has not been explicitly tested, we hypothesise that side branches on the pseudosulci (first- and second-order) facilitate dehydration and subsequent rehydration and together with the spacious ultrastructure facilitate long distance dispersal by air of the pollen grains. A consequence is, however, that pollen with branched pseudosulci (the derived type) germinates slower (KB personal observation, June 2012), perhaps because the pollen first needs to rehydrate on the female structure (or germination medium) before germination can take place. The feature may thus represent a trade-off between dispersal and germination ability. In passing, it is interesting to note that among the many extinct forms of ephedroid pollen, there is a type (common in the Eocene) with extensively branched pseudosulci, and side branches extending almost to the ridge of the plicae (*Ephedripites* (subgenus *Distachyapites*) *claricristatus*). This type, albeit with some modification, is also known from the Neogene, but not from extant species of *Ephedra*.

**Number of plicae.** — The number of plicae is mostly consistent within species although some intraspecific variation occurs (Figure 12A). It is further clear that there has been a trend towards reduction of the number of plicae during the course of evolution in *Ephedra* (Figure 14A). The morphology and structure of the exine are known to have substantial implications for the pollination biology of plants (Ferguson & Skvarla 1982; Grayum 1986; Osborn et al. 1991; Bolinder et al. 2015). For entomophilous


pollen to be successfully transported from the microsporangiate structures to the ovulate structures (specifically in this case, to the pollination drop), the pollen needs to adhere to an insect vector. Insect-pollination is probably the ancestral mode of pollination in the Gnetales (Bolinder et al. 2014; Rydin & Bolinder 2015) and pollen of the entomophilous *Welwitschia* (Pearson 1907; Wetschnig & Depish 1999) and *Ephedra foeminea* is sticky and forms distinct clumps (Hesse 1984; Bolinder et al. 2014, 2015). However, since pollenkit is lacking in the Gnetales (Hesse 1984), the means, by which this stickiness is accomplished is currently unknown. In addition to the observed stickiness (Hesse 1984; Bolinder et al. 2014, 2015), we suggest that the numerous plicae facilitate attachment to the setae on the body of an insect vector. Pollen of *Welwitschia* is similar to the ancestral *Ephedra* pollen type in many respects; it is also polyplcate and ellipsoid with the longest axis equal to one of the equatorial axes (Carafa et al. 1996; El-Ghazaly et al. 1998). But in contrast to that of *Ephedra*, *Welwitschia* pollen has a single broad sulcus extending parallel to the long equatorial axis, where the exine splits open at germination (Rydin & Friis 2005). Furthermore, the area between the plicae differs ultrastructurally between the two genera. In *Welwitschia* grains, both tectum and infratectum are present in the furrows between plicae, but in *Ephedra* grains these layers are absent in the furrow regions (Osborn 2000; Bolinder et al. 2015). The functional implication is that while the furrows of *Ephedra* are zones of weakness that can function as apertures, those of *Welwitschia* are not. The third member of the Gnetales, *Gnetum*, is also thought to be insect-pollinated (Kato & Inoue 1994;

←  
Figure 11. Variation in pollen traits among clades (as defined in Figures 13 and 14) and pollen types. **A.** Number of plicae in the different grades/clades. There are considerably more plicae in pollen of the Mediterranean and South American species than in the Asian and North American species. **B.** Number of plicae in pollen with different pseudosulcus morphology. Pollen with unbranched pseudosulci has significantly more plicae than pollen with branched pseudosulci. Pollen with first- and second-order branched pseudosulci has fewer plicae than pollen with first-order branches only. **C.** Length of the long equatorial axis in different clades. There is no noteworthy difference among pollen of the species in the different grade/clades. **D.** Length of the long equatorial axis in pollen of different pseudosulcus morphology. Pollen with first- and second-order branches on the pseudosulci is slightly larger than pollen with unbranched pseudosulci or pseudosulci with first-order branching only. **E.** P/E ratio in different clades. Pollen of Asian species tends to have a smaller P/E ratio than pollen of remaining species. **F.** P/E ratio in pollen with different pseudosulcus morphology. There is no significant difference in P/E ratio among pollen with different pseudosulcus morphology.

→  
Figure 12. Variation in pollen traits among species. Mediterranean species in red, North American species in blue, South American species in green and Asian species in pink. **A.** Number of plicae. Pollen of Mediterranean and South American species generally has a larger number of plicae than does pollen of North American and Asian species. There is relatively high intraspecific variation in this character and it overlaps extensively among species. **B.** Size as gauged by the length of the long equatorial axis. There is some variation among species but the large intraspecific variation, in particular in the North American and Asian species, makes the overlap extensive among species and clades. **C.** Shape as assessed by the P/E ratio. There is some variation among species but the large intraspecific variation makes the overlap extensive among species and clades.

A

species



B

species




Figure 12. (Continued).

C

species






Figure 12. (Continued).

A



B



Kato et al. 1995). Pollen of this genus is spherical with a large number of spines covering the exine surface (Woodhouse 1935; Osborn 2000; Yao et al. 2004), which probably also facilitates adherence to the bodies of insects. It has further been suggested that the spines of *Gnetum* pollen are homologous with the plicae of *Ephedra* and *Welwitschia* pollen (Osborn 2000), which further supports numerous ‘plicae’ (modified into spines in *Gnetum*) as the ancestral state in the Gnetales.

**Pollen size.** — The size of pollen grains varies tremendously within species of *Ephedra*, and does in many cases not overlap among individuals of a single species. Even though the interspecific variation in size is greater than the intraspecific variation, we do not find the character useful for species identification due to the large overlap among species. For the same reason, it is problematic to assign dispersed fossil ephedroid pollen grains to a species (or even clade) based solely on size. Furthermore, the variation in size is not phylogenetically informative ( $\lambda = 0$ ; Figure 14B) and the evolutionary conclusions that can be drawn from pollen size variation are therefore limited. Also the variation in shape (P/E ratio) is large within and among species. Again, the variation between species is larger than within species (Figure 12C), but there is a lot of overlap and the P/E ratio cannot be considered an informative character. The P/E ratio of North American species does not differ significantly from that of South American or Mediterranean species, indicating that North American species have retained the ancestral shape of pollen grains also present in Mediterranean and South American species. This is further supported by the estimate of phylogenetic signal and ancestral state reconstruction ( $\lambda = 0$ ; Figure 14C). There is a

tendency for species in the Asian clade to have smaller mean P/E ratios, meaning that pollen of Asian species have evolved a different shape compared to that of species in the other clades (Figures 12C, 14C).

**Comparison with previous work.** — The morphology of the pseudosulci has, together with number of plicae, traditionally been used to classify *Ephedra* pollen into 2–4 different pollen types (Woodhouse 1935; Steeves & Barghoorn 1959; Zhang & Xi 1983; Kedves 1987; Freitag & Maier-Stolte 1994). The ancestral type defined here is equivalent to the ‘fragilis type’ described by Beug (1956) and Freitag and Maier-Stolte (1994), and to type D described by Steeves and Barghoorn (1959) and Zhang and Xi (1983). The derived type defined here is equivalent to the ‘distachya type’ described by Beug (1956) and Freitag and Maier-Stolte (1994), and this type was divided into several subgroups (types A, B and C) by Steeves and Barghoorn (1959) and (types A and BC) by Zhang and Xi (1983). None of the previously described pollen types (A, B, C) corresponds to pollen of the Asian or North American species and we find no support for any of these previously suggested delimitations within the derived pollen type. Taken together, our results demonstrate that the taxonomic value of pollen morphology is limited.


**Pollen dimorphism.** — Several kinds of intraspecific pollen dimorphism have been reported. Kedves (1987) reported a dimorphism in size in several *Ephedra* species and argues that variation in size, therefore, is not a valuable taxonomic character. We have not observed size dimorphism in our data. Ickert-Bond et al. (2003) reported dimorphism,

←

Figure 13. Ancestral state reconstruction of pseudosulci branching. **A.** Branches on the pseudosulci coded as a binary character: Absence of side branches (0) and presence of side branches (1). Absence of side branches on the pseudosulci is the ancestral state in *Ephedra*. Presence of branches on the pseudosulci is inferred to have evolved independently twice, once along the stem lineage of the Asian clade and once on along the stem lineage of the North American clade. There are reversals to the ancestral state in both clades. **B.** Side branches on the pseudosulci as a multistate character: Absence of branches (0), presence of first-order branches (1) and presence first- and second-order branches (2). First-order branches on the pseudosulci are inferred to have originated several times. The character has subsequently been lost several times among North American species and once in the Asian clade. Second-order branching of the pseudosulci has evolved at least once in both clades.

→

Figure 14. Ancestral state reconstruction of the mean of number of plicate, size and shape of pollen grains. **A.** Mean number of plicae. During the course of evolution in *Ephedra* there has been a general trend towards reduction of the number of plicae in the North American and Asian clades. Numerous plicae have, however, evolved at least twice among the North American species. Mediterranean and South American species generally have an intermediate number of plicae, and numerous plicae have evolved several times. **B.** Size (mean length of the long equatorial axis). The variation in size in the genus is not phylogenetically informative. **C.** Shape (mean P/E ratio). There is no difference in P/E ratio between Mediterranean, South American and North American species. Pollen of Asian species has, in general, a smaller P/E ratio than that of the other grade/clades.



regarding presence and absence of first-order branches on the pseudosulci, for a probable hybrid between *E. funerea* and *E. torreyana*. This finding is indirectly supported by our results since *E. funerea* has the derived pollen type with side branches on the pseudosulci, whereas *E. torreyana* probably represents a reversal back to the ancestral state and has pollen without side branches. Ickert-Bond et al. (2003) also showed the presence of two types of pollen in the same microsporangium of *E. trifurca* (with and without first-order branches), but this observation is not supported by the data presented here. However, we have seen similar examples in some specimens of *E. alata* that are of probable hybrid origin, as indicated by our results and molecular data (CR personal observation, December 2014). In these specimens, a fraction of the pollen has side branches on the pseudosulci while the majority lacks the feature (as does all pollen in most specimens of *E. alata*; Figures 3B, 7B). Specimens of putative hybrid origin were, however, removed from the present study at an early stage.

The dimorphism described by El-Ghazaly and Rowley (1997), Ickert-Bond et al. (2003) and Doores et al. (2007), concerning a curvature of the ridges (plicae), is not supported by our data. These studies describe a 'normal' pollen form with straight ridges and a variant form with sinuous ridges. We have not observed such dimorphism in any pollen from any accession investigated in the present study. Instead, we have seen that treatment of pollen with alcohol alters the size and shape of *Ephedra* pollen considerably (Figure 2), as does preparation with acetolysis (Kedves 1987), which typically affects some but not all grains. In all studies reporting dimorphism of the ridges (El-Ghazaly & Rowley 1997; Ickert-Bond et al. 2003; Doores et al. 2007), pollen was treated with alcohol, acetolysed and/or critical-point dried, and reported intraspecific dimorphism concerning sinuous or straight ridges is most likely a consequence of specimen preparation.

## Conclusions and evolutionary implications

Our results show that *Ephedra* pollen occurs in two distinct forms, an ancestral type (with unbranched pseudosulci) present in Mediterranean and South American species, and a derived type (with branched pseudosulci) that appears to have evolved independently twice, once in the Asian clade and once in the clade containing all North American species except *E. pedunculata*. There are repeated reversals back to the ancestral state within both these clades. Although there is phylogenetic information in several pollen features, the repeated reversals in combination with some degree of intraspecific variation make it diffi-

cult to assign individual pollen grains to species or subclades of the genus. Further, we find no clear correlation between the two pollen types and pollination syndrome. Species that have the ancestral type may be insect-pollinated (such as *E. foeminea*; Bolinder et al. 2014, 2015) or wind-pollinated (such as *E. trifurca*; e.g. Niklas et al. 1986). The derived pollen type is, however, only known from wind-pollinated species.

Polyaplicate pollen is not unique to the Gnetales; it also occurs in several extant and extinct angiosperms, such as the Alismatales, Laurales and Zingiberales (Hesse et al. 2000; Friis et al. 2004). Furthermore, it is far from clear that all dispersed polyaplicate pollen in Mesozoic strata referred to as 'ephedroids', was produced by a single group of plants. However, the gnetalean affinity of individual grains can be assessed based on ultrastructural studies of the pollen wall (Hesse et al. 2000; Tekleva & Krassilov 2009; Friis et al. 2011). Ultrastructural information is only available for a few Mesozoic ephedroids (Trevisan 1980; Osborn et al. 1993; Kedves 1994), and those are clearly of the ancestral ephedroid pollen type described here (see also Bolinder et al. 2015). Germinated pollen grains (i.e. shed exines) are also found *in situ* in *Ephedra* seeds from the Early Cretaceous (Rydin et al. 2004, 2006a) and, as assessed by their many plicae and the absence of side branches on the pseudosulci, this pollen is also clearly of the ancestral type. The same inference can be made for other dispersed ephedroid pollen grains found in Mesozoic strata (although the ultrastructure has not been studied for any of these grains) (Srivastava 1968; Scott 1960; Wilson 1962; Stover 1964; Muller 1968; Brenner 1976; de Lima 1980; Osborn et al. 1993; Takahashi 1995; Narváez & Sabino 2008; Abubakar et al. 2011).

Ephedroid pollen with branched pseudosulci (i.e. the derived type) has, to our knowledge, only been described twice from the Mesozoic: once from the lower Upper Cretaceous Raritan Formation in North America (Steeves & Barghoorn 1959) and once from the Xining Basin of the Tibetan Plateau (Norbäck Ivarsson 2014), found in a section from the Cenomanian–Maastrichtian (Horton et al. 2004). After the K-Pg boundary, pollen of the derived type gradually becomes much more common and dominates over the ancestral type in most palaeo-palynofloras from the Paleocene and onwards (Cookson 1956; Gray 1960; Ghosh et al. 1963; Nagy 1963; Shaw 1998; Hoorn et al. 2012). This increase of the derived pollen type in the Cenozoic probably represents an adaptation to climatic changes after the K-Pg boundary. Previous studies have indicated an early Oligocene (Ickert-Bond et al. 2009) or even younger (Huang & Price 2003) age of

crown-group *Ephedra*. The results of the present study and a first occurrence of fossil pollen of the derived type in the Late Cretaceous indicate that the crown-group *Ephedra* is much older than previously estimated.

### Acknowledgements

The authors thank the directors of the herbaria A, ASU, B, BM, BR, CAI, E, F, GB, GH, L, LL, M, MO, MSB, NMC, NY, KAS, PE, RAS, S, TEX, UPS, W and WU for access to plant material, and Wafaa M. Amer (Cairo University), Julien B. Bachelier (State University of New York), Chen Hou and Eva Larsén (Stockholm University) for selecting appropriate specimens, Kjell Jansson (Stockholm University) for laboratory assistance, Reinhard Zetter and Friðgeir Grimsson (University of Vienna) for fruitful discussions and Brian K. Horton (University of Texas at Austin) for sharing fossil material. The authors also wish to thank James A. Doyle and one anonymous reviewer for constructive insights and helpful comments on the paper.

### Disclosure statement

No potential conflict of interest was reported by the authors.

### Funding

Funding was provided from Stiftelsen Lars Hiertas Minne (to KB) and from the Swedish Research Council VR (to CR).

### Specimens investigated

*Mediterranean species.* – *Ephedra alata* Decne., KH Rechinger 102 (S), AA Anderberg 480 (S), Ibrahim & Mahdi s.n. (S), AbdEaleem s.n. (CAI); *E. altissima* Desf., A Faure s.n. (S), M Staudinger 6714 (W), H Freitag 35035 (KAS); *E. aphylla* Forssk., G Samuelsson 2696 (S), F Wettstein 2751 (WU), J Bornmüller 1746 (WU), WM Amer (CAI); *E. ciliata* Fisch. et C.A.Mey., NT Yakovleva (W), EK Balls 2487 (S), H Handel-Mazetti 973 (WU), JET Aitchison 496537 (GH); *E. foeminea* Forssk., K Bolinder 542 (S), L Norbäck Ivarsson & O Thureborn s.n. (S), R Pamaninio & R Pichisermolli 139 (L), FS Meyers & JE Dinsmoore 8124 (L), JS Andersen et al. 2098 (S), Bornmüller 1746 (A), O Porch s.n. (WU); *E. foliata* Boiss et C.A.Mey., B Tiagi s.n. (S), Wendelbo & Assadi 16604 (W), Hedberg & Hedberg 92019A V-60888 (UPS), Thulin & Al-Gifri 9975 (V-096087) 183895 (UPS), PN Parker 3304 (A); *E. milleri* Freitag et Maier-St., P Hein et al. YP 1110 (B), Miller 7667A (E).

*North American species.* – *Ephedra antisiphilitica* Berlandier ex C.A.Mey., HC Hanson 344 (NY), ME Jones 3726 (BR), Hendersoin 62-02a (BR), E Palmer 1292 (GH); *E. aspera* Engelm.,

KF Parker 7799 (NY), NH Holmgren 6604 (NY), RD Worthington 13620 (NY), BA Stein 31 (RSA); *E. californica* S.Watson, FM Reed 5772 (L), C Epling & WM Robinson s.n. (L), A Carter 3667 (L); *E. compacta* Rose, DS Corell & IM Johnston 20233 (NY); *E. coryi* E.L. Reed, SM Ickert-Bond 953 (ASU), DS Corell 32805 (S), DS Corell 32785 (S); *E. cutleri* Peebles, CT Mason Jr. 2192 (ASU), NH Holmgren 12744 (NY), DE Atha et al. s.n. (NY); *E. fasciculata* A.Nelson, ME Jones s.n. (RSA), FW Gould 1526 (GH), JH Lehr 2309 (NY), PA Munz 12053 (RSA); *E. funerea* Coville et C.V.Morton, J Wash & IW Clokey 8224 (NY), LS Rose 67021 (S), CL Hitchcock 12329 (A); *E. nevadensis* S.Watson, IW Clokey 6509 (S), C Epling & W Robinson (S), P Raven 14251 (WU), LS Rose 58108 (NY), *E. pedunculata* Engelm. ex S.Watson, Johnston 8847 (TEX, LL), Hendrickson 23183 (TEX, LL), RM Steward 2265 (GH); *E. torreyana* S.Watson, RW Spellenberg 10204 (NMC), IW Clokey 7816 (NY), E Neese & K Mutz 11414 (NY), Johnston et al. 10578 (F); *E. trifurca* Torr., RD Worthington 24587 (NY), A Nelson 1619 (NY), MC Johnston et al. 10573 (NY), P Allen (S), CV Hartman 642 (GH); *E. viridis* Coville, LS Rose 58080 (S), JL Reveal 100 (NY), Neely 4353 (NY), JT Howell 3824 (GH).

*South American clade.* – *Ephedra americana* Humb., Bonpl. ex Willd., Gerth s.n. (L), E Günter & O Buchtien (S), RE Fries 1044 (S), M Cardenas 4 (GH); *E. boelckeii* F.A.Roig, Maas et al. 8184 (GB); *E. breana* Phil., KH Rechinger & W Rechinger 63547 (W), IM Johnston 3613 (GH); *E. chilensis* C.Presl., C Skottsberg 987 (F), Werdermann 138 (E), Jouffé 2751 (GH); *E. frustillata* Miers, MP Moreno 236 (NY), JB Hatcher s.n. (NY), LR Parodi 11858 (GH); *E. multiflora* Phil. ex Stapf, Baines et al. 233 (E), J Chiapella 2344 & E Vitek 09-0359 (W), Johnson 6286 (GH); *E. ochreatea* Miers, A Donat 34 (GB), MP Moreno 433 (NY), RH Fortunato 5413 (NY), W Fischer 15 (GH); *E. triandra* Tul., J Chiapella 2505 & E Vitek 09-0520 (W), A Cuzzo 1954 (S), K Fiebrig 215 (GH); *E. trifurcata* Zöllner, O Zöllner 7928 (L); *E. tweediana* Fisch. et C.A.Mey., R Gallinal 5683 (GH), C Osten 22008 (S), S Venturi 348 (S), JH Hunziker 1648 (S).

*Asian clade.* – *Ephedra distachya* L., J Prudhomme 89 (WU), K Bolinder 764 (S), I Rácz et al 35308 (S), Rechinger 53066 (W), Flora Germanica Exsiccatae 2325 (GH); *E. equisetina* Bunge, RC Ching 109 (MO), QR Wu s.n. (MO), KK Andrjoschenko 4027 (S), E Mokeeva (A); *E. gerardiana* Wall. ex Florin, P Wendelbo s.n. (WU), Walter Koelz 5310 (S), P Wendelbo s. n. (S), Österreichische Karako Exp. 1039 (W), Stainton, Syes & Williams 813 (E); *E. intermedia* Schrenk et C.A.Mey., Lindberg 117-1947 (W), W Koeltz 2305e (NY), Y Yang 08070801 (PE), JET Aitchison 1122 (GH); *E. likiangensis* Florin, JF Rock 3694 (NY), G Forrest 5564 (BR); *E. lomatolepia* Schrenk, II Rusanovick & LA Krai 56 (NY); *E. minuta* Florin, B Dichore 8457 (MSB), Long et al. 153 (W); *E. monosperma* J.G.Gmel. ex C.A. Mey., H Freitag 33068 (KAS), Potanin s.n. (BR), CG 81-0152 (BR); *E. pachyclada* Boiss., KH & F Rechinger 3676 (W), A Danin S-2455 (S), JP Mandaville s.n. (BM); *E. regeliana* Florin, H Hastman 26 (MSB), GL Webster & E Nasir 5959 (W), GL Webster & E Nasir 5950 (GH); *E. sarcocarpa* Aitch. et Hemsl., F Rechinger 46054b (W), Ruttner 594 (W), Tsanshakt & Riedel 15977 (W), F Rechinger 46295 (W), Aellen & Estafandri 2786 C-7840 (S); *E. saxatilis* (Stapf) Royle ex Florin, F Lobblicher 87 (M), CE Parkinson 7077 (S); *E. sinica* Stapf, E Licent 13523 (S), Y Yang 99531 (PE), W Qingru 9812 (MO); E Licewtak 13523 (A); *E. strobilacea* Bunge, Supra 3369a (NY), A Michelson 3369b (S), KH Rechinger & F Rechinger 2703 (S), Leonard 5582 (BR); *E. transitoria* Riedl, V Täckholm et al. 9166 (BR), Doppelbaur 190 (M), Rechinger 8990 (W).

Table I. Pollen properties and distribution for all investigated *Ephedra* species.

| Clade, species                                    | Distribution                             | Polar axis ( $\mu\text{m}$ ) | Equatorial diameter ( $\mu\text{m}$ ) | P/E ratio                   | Number of plicae       | Pseudosulci                                                    |
|---------------------------------------------------|------------------------------------------|------------------------------|---------------------------------------|-----------------------------|------------------------|----------------------------------------------------------------|
| <i>Mediterranean species</i>                      |                                          |                              |                                       |                             |                        |                                                                |
| <i>E. foeminea</i> Forssk.                        | Eastern Mediterranean                    | 18.1 (13.6–23.6) $\pm$ 2.3   | 42.2 (33.8–53.1) $\pm$ 3.8            | 0.43 (0.34–0.55) $\pm$ 0.05 | 15.8 (12–22) $\pm$ 2.6 | Not branched                                                   |
| <i>E. alata</i> Decne.                            | North Africa; Near East                  | 20.0 (16.7–23.2) $\pm$ 1.6   | 58.4 (52.5–65.4) $\pm$ 3.2            | 0.34 (0.28–0.43) $\pm$ 0.03 | 14.9 (12–16) $\pm$ 1.5 | Not branched                                                   |
| <i>E. altissima</i> Desf.                         | North Africa                             | 17.5 (14.5–20.9) $\pm$ 1.5   | 42.5 (38.0–50.6) $\pm$ 3.2            | 0.41 (0.32–0.50) $\pm$ 0.05 | 16.5 (12–20) $\pm$ 1.6 | Not branched                                                   |
| <i>E. aphylla</i> Forssk.                         | North Africa; Near East                  | 21.6 (16.8–28.0) $\pm$ 2.2   | 49.4 (38.3–58.0) $\pm$ 3.7            | 0.44 (0.32–0.63) $\pm$ 0.06 | 15.5 (10–20) $\pm$ 2.2 | Not branched                                                   |
| <i>E. ciliata</i> Fisch. et C.A.Mey.              | Mediterranean to Central Asia            | 17.6 (14.2–22.8) $\pm$ 1.7   | 40.2 (33.0–46.4) $\pm$ 2.7            | 0.44 (0.35–0.59) $\pm$ 0.05 | 17.5 (10–22) $\pm$ 2.7 | Not branched                                                   |
| <i>E. foliata</i> Boiss. et C.A.Mey.              | Southeast Mediterranean to Central Asia  | 17.6 (13.3–22.8) $\pm$ 1.8   | 41.3 (33.6–51.2) $\pm$ 3.4            | 0.42 (0.29–0.55) $\pm$ 0.04 | 16.7 (10–22) $\pm$ 2.7 | Not branched                                                   |
| <i>E. milleri</i> Freitag et Maier-St.            | Endemic to Oman                          | 20.5 (16.1–24.4) $\pm$ 1.9   | 46.8 (38.1–57.0) $\pm$ 3.5            | 0.44 (0.36–0.52) $\pm$ 0.04 | 16.3 (10–20) $\pm$ 2.0 | Not branched                                                   |
| <i>North American (including Mexico) species</i>  |                                          |                              |                                       |                             |                        |                                                                |
| <i>E. antisiphilitica</i> Berlandier ex. C.A.Mey. | Southwest North America; North Mexico    | 19.4 (9.1–28.3) $\pm$ 4.5    | 40.2 (20.2–55.34) $\pm$ 11.4          | 0.49 (0.37–0.69) $\pm$ 0.07 | 16.6 (12–22) $\pm$ 2.7 | Rarely short first-order branches, no secondary branches       |
| <i>E. aspera</i> Engelm.                          | Southwest North America; North Mexico    | 17.7 (8.0–28.5) $\pm$ 5.5    | 39.5 (19.5–58.1) $\pm$ 13.8           | 0.46 (0.34–0.61) $\pm$ 0.06 | 13.0 (10–18) $\pm$ 2.0 | Short first-order branches, no secondary branches              |
| <i>E. californica</i> S. Watson                   | California; Mexico Baja                  | 18.5 (9.4–29.1) $\pm$ 6.3    | 46.2 (26.3–64.0) $\pm$ 13.2           | 0.39 (0.29–0.55) $\pm$ 0.05 | 14.5 (10–20) $\pm$ 2.3 | Not branched                                                   |
| <i>E. compacta</i> Rose                           | Central Mexico                           | 17.3 (13.9–20.7) $\pm$ 1.7   | 37.3 (31.7–43.7) $\pm$ 2.9            | 0.46 (0.37–0.55) $\pm$ 0.05 | 12.5 (10–16) $\pm$ 1.7 | Occasionally short first-order branches, no secondary branches |
| <i>E. coryi</i> E.L.Reed                          | New Mexico, Texas                        | 14.9 (9.8–28.1) $\pm$ 6.0    | 33.9 (20.7–59.3) $\pm$ 13.4           | 0.44 (0.31–0.52) $\pm$ 0.04 | 6.5 (4–10) $\pm$ 1.5   | Long first-order branches, often secondary branched            |
| <i>E. cutleri</i> Peebles                         | Arizona, Colorado, New Mexico, Utah      | 19.4 (8.6–26.1) $\pm$ 4.3    | 47.9 (24.3–58.9) $\pm$ 9.6            | 0.41 (0.28–0.49) $\pm$ 0.04 | 7.0 (4–10) $\pm$ 1.3   | First-order branches, occasionally secondary branched          |
| <i>E. fasciculata</i> A. Nelson                   | Arizona, California, Nevada, Utah        | 20.0 (9.3–30.6) $\pm$ 6.1    | 46.7 (22.2–63.8) $\pm$ 13.4           | 0.43 (0.33–0.53) $\pm$ 0.05 | 10.1 (8–12) $\pm$ 1.1  | First-order branches, no secondary branches                    |
| <i>E. funerea</i> Coville et C.V.Morton           | California to Nevada (Death Valley area) | 10.9 (9.3–12.2) $\pm$ 0.65   | 27.4 (24.9–30.4) $\pm$ 1.13           | 0.40 (0.32–0.46) $\pm$ 0.03 | 7.2 (5–10) $\pm$ 1.2   | First-order branches, no secondary branches                    |
| <i>E. nevadensis</i> S.Watson                     | Southwest North America                  | 21.9 (11.4–30.7) $\pm$ 5.5   | 52.8 (24.7–73.3) $\pm$ 14.2           | 0.42 (0.29–0.59) $\pm$ 0.07 | 7.7 (6–12) $\pm$ 1.4   | First-order branches, occasionally secondary branched          |
| <i>E. pedunculata</i> Engelm. ex S.Watson         | Texas, North Mexico                      | 18.4 (13.6–23.7) $\pm$ 1.9   | 42.6 (33.6–53.5) $\pm$ 4.6            | 0.43 (0.30–0.59) $\pm$ 0.06 | 15.0 (12–18) $\pm$ 1.4 | Not branched                                                   |
| <i>E. torreyana</i> S.Watson                      | Southwest North America; North Mexico    | 17.9 (14.2–22.9) $\pm$ 1.4   | 42.3 (28.2–51.8) $\pm$ 6.3            | 0.43 (0.29–0.64) $\pm$ 0.07 | 14.5 (10–20) $\pm$ 2.1 | Not branched                                                   |
| <i>E. trifurca</i> Torr. Ex S.Watson              | Southwest North America; North Mexico    | 18.4 (11.7–24.7) $\pm$ 2.4   | 44.1 (35.5–54.7) $\pm$ 5.1            | 0.44 (0.31–0.64) $\pm$ 0.08 | 16.2 (10–22) $\pm$ 2.5 | Not branched                                                   |

|                                               |                                    |                        |                          |                         |                    |                                                                |
|-----------------------------------------------|------------------------------------|------------------------|--------------------------|-------------------------|--------------------|----------------------------------------------------------------|
| <i>E. viridis</i> Coville                     | Southwest North America            | 24.4 (18.4–30.4) ± 2.7 | 54.1 (42.2–65.7) ± 5.2   | 0.45 (0.30–0.57) ± 0.05 | 6.9 (4–10) ± 1.2   | First-order branches, occasionally secondary branched          |
| <i>South American species</i>                 |                                    |                        |                          |                         |                    |                                                                |
| <i>E. americana</i> Humb. et Bonpl. ex Willd. | Andean South America + Argentina   | 18.5 (14.3–24.9) ± 2.1 | 42.8 (31.7–64.0) ± 7.1   | 0.44 (0.31–0.61) ± 0.06 | 13.9 (10–18) ± 1.7 | Not branched                                                   |
| <i>E. boelckei</i> F.A.Roig                   | Argentina                          | 15.8 (13.4–17.7) ± 1.0 | 38.0 (34.1–42.3) ± 2.0   | 0.42 (0.35–0.47) ± 0.03 | 13.5 (10–16) ± 1.4 | Not branched                                                   |
| <i>E. breana</i> Phil.                        | Argentina & Chile                  | 20.9 (16.5–25.6) ± 2.5 | 45.3 (39.7–50.0) ± 2.8   | 0.46 (0.37–0.63) ± 0.07 | 16.3 (12–20) ± 2.2 | Not branched                                                   |
| <i>E. chilensis</i> C.Presl.                  | Andean South America               | 20.7 (17.9–24.1) ± 1.2 | 45.0 (40.2–51.7) ± 2.4   | 0.46 (0.40–0.56) ± 0.03 | 14.2 (10–20) ± 2.3 | Not branched                                                   |
| <i>E. frustillata</i> Miers                   | South Argentina; Chile             | 17.6 (12.7–23.9) ± 2.4 | 40.6 (35.4–47.7) ± 2.6   | 0.43 (0.32–0.60) ± 0.06 | 13.5 (10–18) ± 1.8 | Not branched                                                   |
| <i>E. multiflora</i> Phil. ex. Stapf          | North Chile; Argentina             | 17.4 (13.5–22.0) ± 2.0 | 43.2 (35.7–50.7) ± 3.6   | 0.40 (0.32–0.49) ± 0.03 | 13.6 (10–18) ± 2.0 | Not branched                                                   |
| <i>E. ochreatea</i> Miers                     | Argentina                          | 16.9 (12.4–26.1) ± 3.3 | 40.4 (31.9–50.6) ± 4.7   | 0.42 (0.33–0.57) ± 0.05 | 15.1 (12–20) ± 1.9 | Not branched                                                   |
| <i>E. triandra</i> Tul.                       | Argentina to South Brazil          | 18.2 (15.1–22.2) ± 1.6 | 35.8 (25.7–49.2) ± 7.7   | 0.53 (0.33–0.74) ± 0.09 | 18.8 (14–24) ± 2.6 | Not branched                                                   |
| <i>E. trifurcata</i> Zöllner                  | Chile                              | 9.8 (8.0–11.3) ± 0.7   | 20.5 (18.2–23.04) ± 1.22 | 0.48 (0.41–0.53) ± 0.03 | 12.3 (10–14) ± 1.0 | Not branched                                                   |
| <i>E. tweediana</i> Fisch. et C.A.Mey.        | Uruguay and Argentina              | 15.6 (12.2–19.8) ± 1.6 | 37.6 (30.6–44.2) ± 3.0   | 0.41 (0.32–0.62) ± 0.06 | 14.2 (10–20) ± 1.8 | Not branched                                                   |
| <i>Asian species</i>                          |                                    |                        |                          |                         |                    |                                                                |
| <i>E. distachya</i> L.                        | Europe and Central Asia            | 20.0 (15.0–25.1) ± 1.8 | 49.6 (41.7–59.9) ± 3.7   | 0.41 (0.32–0.53) ± 0.04 | 6.6 (4–10) ± 1.2   | First-order branches, occasionally secondary branched          |
| <i>E. equisetina</i> Bunge                    | Central Asia; Russia; North China  | 17.3 (6.4–28.8) ± 6.8  | 38.5 (19.5–56.7) ± 13.8  | 0.44 (0.31–0.60) ± 0.06 | 5.5 (4–8) ± 1.1    | First-order branches, often secondary branched                 |
| <i>E. gerardiana</i> Wall. ex Florin          | Himalaya region to southwest China | 19.7 (15.1–24.4) ± 1.8 | 45.7 (39.2–54.8) ± 2.9   | 0.43 (0.33–0.55) ± 0.05 | 6.4 (4–10) ± 1.3   | First-order branches, no secondary branches                    |
| <i>E. intermedia</i> Schrenk et C.A.Mey.      | Central Asia                       | 15.1 (8.7–24.0) ± 4.9  | 37.0 (20.2–52.2) ± 11.4  | 0.41 (0.30–0.52) ± 0.04 | 9.2 (6–14) ± 1.5   | First-order branches, no secondary branches                    |
| <i>E. tibiangensis</i> Florin                 | Southwest China                    | 20.3 (15.1–27.1) ± 2.4 | 46.9 (35.4–56.0) ± 5.0   | 0.44 (0.30–0.59) ± 0.06 | 13.0 (10–18) ± 2.2 | Occasionally short first-order branches, no secondary branches |
| <i>E. lonatolepis</i> Schrenk                 | Central Asia                       | 12.6 (7.7–21.7) ± 5.9  | 32.3 (20.3–51.7) ± 12.8  | 0.38 (0.32–0.47) ± 0.04 | 11.5 (10–14) ± 1.3 | Not branched                                                   |
| <i>E. minuta</i> Florin                       | West China                         | 15.9 (7.7–25.5) ± 6.6  | 35.4 (20.4–53.1) ± 13.4  | 0.45 (0.35–0.56) ± 0.05 | 15.2 (12–20) ± 2.4 | Occasionally short first-order branches, no secondary branches |
| <i>E. monosperma</i> J.G.Gmel. ex C.A.Mey.    | Siberia to northwest China         | 15.4 (8.2–21.9) ± 5.1  | 36.1 (19.9–53.2) ± 10.9  | 0.42 (0.33–0.53) ± 0.05 | 9.8 (6–12) ± 1.6   | First-order branches, no secondary branches                    |

(Continued)

Table 1. (Continued).

| Clade, species                              | Distribution                | Polar axis ( $\mu\text{m}$ ) | Equatorial diameter ( $\mu\text{m}$ ) | P/E ratio                   | Number of plicae       | Pseudosulci                                           |
|---------------------------------------------|-----------------------------|------------------------------|---------------------------------------|-----------------------------|------------------------|-------------------------------------------------------|
| <i>E. pachyclada</i> Boiss.                 | Iran to Afghanistan         | 18.0 (8.8–25.32) $\pm$ 5.1   | 46.0 (20.6–59.1) $\pm$ 13.8           | 0.39 (0.32–0.50) $\pm$ 0.04 | 6.7 (4–8) $\pm$ 1.2    | First-order branches, occasionally secondary branched |
| <i>E. regeliana</i> Florin                  | Central to East Asia        | 13.3 (7.5–18.2) $\pm$ 3.4    | 36.4 (19.2–46.7) $\pm$ 9.5            | 0.37 (0.29–0.44) $\pm$ 0.03 | 9.4 (6–12) $\pm$ 1.3   | Short first-order branches, no secondary branches     |
| <i>E. sarocarpa</i> Aitch. et Hemsl.        | Afghanistan; Pakistan       | 12.0 (7.0–19.7) $\pm$ 4.4    | 31.3 (19.1–48.6) $\pm$ 10.5           | 0.38 (0.28–0.50) $\pm$ 0.04 | 14.4 (10–20) $\pm$ 2.9 | First-order branches, no secondary branches           |
| <i>E. saxatilis</i> (Stapf) Royle ex Florin | West China                  | 18.8 (8.4–23.7) $\pm$ 3.9    | 46.0 (19.8–57.0) $\pm$ 9.9            | 0.41 (0.33–0.52) $\pm$ 0.04 | 10.9 (8–18) $\pm$ 2.4  | First-order branches, no secondary branches           |
| <i>E. sinica</i> Stapf                      | North China; Mongolia       | 19.0 (8.1–25.6) $\pm$ 4.4    | 47.5 (21.2–62.6) $\pm$ 10.7           | 0.40 (0.26–0.52) $\pm$ 0.05 | 6.8 (4–10) $\pm$ 1.4   | First-order branches, often secondary branched        |
| <i>E. snobilacea</i> Bunge                  | Central Asia to Afghanistan | 15.6 (7.1–26.91) $\pm$ 5.2   | 39.1 (19.6–58.5) $\pm$ 12.0           | 0.39 (0.32–0.56) $\pm$ 0.04 | 13.4 (10–16) $\pm$ 1.6 | Not branched                                          |
| <i>E. transitoria</i> Riedl                 | West Asia                   | 17.1 (9.2–26.2) $\pm$ 4.5    | 40.4 (16.6–52.7) $\pm$ 12.5           | 0.44 (0.31–0.63) $\pm$ 0.08 | 14.3 (10–20) $\pm$ 2.1 | Not branched                                          |

## References

- Abubakar MB, Luterbacher HP, Ashraf AR, Ziedner R, Maigari AS. 2011. Late Cretaceous palynostratigraphy in the Gongola Basin (Upper Benue Trough, Nigeria). *Journal of African Earth Sciences* 60: 19–27. doi:10.1016/j.jafrearsci.2011.01.007.
- Akaike H. 1974. A new look at the statistical model identification. *IEEE Transactions on Automatic Control* 19: 716–723. doi:10.1109/TAC.1974.1100705.
- Akkiraz MS, Kayseri MS, Akgün F. 2008. Palaeoecology of coal-bearing Eocene sediments in Central Anatolia (Turkey) based on quantitative palynological data. *Turkish Journal of Earth Sciences* 17: 317–360.
- Beug H-J. 1956. Pollendimorphismus bei *Ephedra*. *Die Naturwissenschaften* 43: 332–333. doi:10.1007/BF00629402.
- Bharadwaj D. 1963. Pollen grains of *Ephedra* and *Welwitschia* and their probable fossil relatives. *Memoirs of the Indian Botanical Society* 4: 125–135.
- Bino RJ, Dafni A, Meeuse ADJ. 1984. Entomophily in the dioecious gymnosperm *Ephedra aphylla* Forsk. (= *E. alte* C.A. Mey.), with some notes on *E. campylopoda* C.A. Mey. I. Aspects of the entomophilous syndrome. *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen. Series C: Biological and medical sciences* 87: 1–13.
- Bolinder K, Humphreys AM, Ehrlén J, Alexandersson R, Ickert-Bond SM, Rydin C. 2014. Pollination mechanisms in the ancient gymnosperm clade *Ephedra* (Gnetales). In: Bolinder K, ed. *Pollination in Ephedra* (Gnetales), 29–45. Licentiate thesis. Stockholm: Stockholm University.
- Bolinder K, Niklas KJ, Rydin C. 2015. Aerodynamics and pollen ultrastructure in *Ephedra* (Gnetales). *American Journal of Botany* 102: 457–470. doi:10.3732/ajb.1400517.
- Brenner GJ. 1976. Middle Cretaceous floral provinces and early migration of angiosperms. In: Beck CB, ed. *Origin and early evolution of angiosperms*, 23–47. New York: Columbia University Press.
- Buchmann SL, O'Rourke MK, Niklas KJ. 1989. Aerodynamics of *Ephedra trifurca*. III. Selective pollen capture by pollination droplets. *Botanical Gazette* 150: 122–131.
- Carafa AM, Ponzi R, Pizzolongo F. 1996. Some aspects of microsporogenesis of *Welwitschia mirabilis* Hook. *Plant Biosystems* 130: 557–565.
- Cookson IC. 1956. Pollen grains of the *Ephedra* type in Australian tertiary deposits. *Nature* 177: 47–48. doi:10.1038/177047a0.
- Crane PR, Lidgard S. 1989. Angiosperm diversification and paleolatitudinal gradients in Cretaceous floristic diversity. *Science* 246: 675–678. doi:10.1126/science.246.4930.675.
- de Lima MR. 1980. Palinologia da Formação Santana (Cretáceo do Nordeste do Brasil). III. Descrição sistemática dos pólenes da Turma Plicates (Subturma Costates). *Ameghiniana* 17: 15–47.
- Doores AS, Osborn JM, El-Ghazaly G. 2007. Pollen ontogeny in *Ephedra americana* (Gnetales). *International Journal of Plant Sciences* 168: 985–997. doi:10.1086/509538.
- Doyle JA, Endress PK. 2000. Morphological phylogenetic analysis of basal angiosperms: Comparison and combination with molecular data. *International Journal of Plant Sciences* 161: S121–S153. doi:10.1086/ijps.2000.161.issue-s6.
- Doyle JA, Le Thomas A. 1994. Cladistic analysis and pollen evolution in Annonaceae. *Acta Botanica Gallica* 141: 149–170. doi:10.1080/12538078.1994.10515148.
- El-Ghazaly G, Rowley JR. 1997. Pollen wall of *Ephedra foliata*. *Palynology* 21: 7–18. doi:10.1080/01916122.1997.9989483.
- El-Ghazaly G, Rowley JR, Hesse M. 1998. Polarity, aperture condition and germination in pollen grains of *Ephedra* (Gnetales). *Plant Systematics and Evolution* 213: 217–231. doi:10.1007/BF00985202.

- Erdtman G. 1952. Pollen morphology and plant taxonomy: Angiosperms (An introduction to palynology). Stockholm: Almquist & Wiksell.
- Ferguson IK, Skvarla JJ. 1982. Pollen morphology in relation to pollinators in Papilionoideae (Leguminosae). Botanical Journal of the Linnean Society 84: 183–193. doi:10.1111/boj.1982.84.issue-3.
- Franchi G, Nepi M, Dafni A, Pacini E. 2002. Partially hydrated pollen: Taxonomic distribution, ecological and evolutionary significance. Plant Systematics and Evolution 234: 211–227. doi:10.1007/s00606-002-0221-1.
- Freckleton RP, Harvey PH, Pagel M. 2002. Phylogenetic analysis and comparative data: A test and review of evidence. The American Naturalist 160: 712–726. doi:10.1086/an.2002.160.issue-6.
- Freitag H, Maier-Stolte M. 1994. Characterization of areas, Ephedraceae. In: Browicz K, ed. Chorology of trees and shrubs in South-West Asia and adjacent regions, Vol. 10, 5–16. Kornik: Polish Academy of Sciences, Institute of Dendrology.
- Friis EM, Crane PR, Pedersen KR. 2011. Early flowers and angiosperm evolution. New York: Cambridge University Press.
- Friis EM, Pedersen KR, Crane PR. 2004. Araceae from the early Cretaceous of Portugal: Evidence on the emergence of monocotyledons. Proceedings of the National Academy of Sciences, USA 101: 16565–16570. doi:10.1073/pnas.0407174101.
- Ghosh A, Banerji D, Srivastava S. 1963. Ephedra-type pollen grains in the Dharamsala (Tertiary) formations, Punjab, India. Nature 198: 408. doi:10.1038/198408a0.
- Gray J. 1960. Temperate pollen genera in the Eocene (Claiborne) flora, Alabama. Science 132: 808–810. doi:10.1126/science.132.3430.808.
- Grayum MH. 1986. Correlations between pollination biology and pollen morphology in the Araceae, with some implications for angiosperm evolution. In: Blackmore S, Ferguson IK, eds. Pollen and spores: Form and function, 313–327. Orlando: Academic Press Inc.
- Gullvåg BM. 1966. The fine structure of some gymnosperm pollen walls. Grana 6: 435–475.
- Harvey PH, Pagel MD. 1991. The comparative method in evolutionary biology. Oxford: Oxford University Press.
- Hesse M. 1984. Pollenkitt is lacking in Gnetales: Ephedra and Welwitschia; further proof for its restriction to the angiosperms. Plant Systematics and Evolution 144: 9–16. doi:10.1007/BF00990796.
- Hesse M, Weber M, Halbritter H. 2000. A comparative study of the polyplicate pollen types in Arales, Laurales, Zingiberales and Gnetales. In: Harley MM, Morton CM, Blackmore S, eds. Pollen and spores: Morphology and biology, 227–239. Kew: Royal Botanic Gardens.
- Hoorn C, Straathof J, Abels HA, Xu YD, Utescher T, Dupont-Nivet G. 2012. A late Eocene palynological record of climate change and Tibetan Plateau uplift (Xining Basin, China). Palaeogeography, Palaeoclimatology, Palaeoecology 344–345: 16–38. doi:10.1016/j.palaeo.2012.05.011.
- Horton BK, Dupont-Nivet G, Zhou J, Waanders GL, Butler RF, Wang J. 2004. Mesozoic–Cenozoic evolution of the Xining–Minhe and Dangchang basins, northeastern Tibetan Plateau: Magnetostratigraphic and biostratigraphic results. Journal of Geophysical Research 109: 1–15. doi:10.1029/2003JB002913.
- Huang J, Price RA. 2003. Estimation of the age of extant Ephedra using chloroplast *rbcl* sequence data. Molecular Biology and Evolution 20: 435–440. doi:10.1093/molbev/msg049.
- Huynh K-L. 1975. Le problème de la polarité du pollen d'Ephedra. Pollen et Spores 16: 469–474.
- Ickert-Bond SM, Rydin C, Renner SS. 2009. A fossil-calibrated relaxed clock for Ephedra indicates an Oligocene age for the divergence of Asian and New World clades and Miocene dispersal into South America. Journal of Systematics and Evolution 47: 444–456. doi:10.1111/jse.2009.47.issue-5.
- Ickert-Bond SM, Skvarla JJ, Chisoe WF. 2003. Pollen dimorphism in Ephedra L. (Ephedraceae). Review of Palaeobotany and Palynology 124: 325–334. doi:10.1016/S0034-6667(03)00002-2.
- Ickert-Bond SM, Wojciechowski MF. 2004. Phylogenetic relationships in Ephedra (Gnetales): Evidence from nuclear and chloroplast DNA sequence data. Systematic Botany 29: 834–849. doi:10.1600/0363644042451143.
- Kato M, Inoue T. 1994. Origin of insect pollination. Nature 368: 195. doi:10.1038/368195a0.
- Kato M, Inoue T, Nagamitsu T. 1995. Pollination biology of Gnetum (Gnetaceae) in a lowland mixed dipterocarp forest in Sarawak. American Journal of Botany 82: 862–868.
- Kedves M. 1987. LM and EM studies on pollen grains of recent Welwitschia mirabilis Hook. and Ephedra species. Acta Botanica Hungarica 33: 81–103.
- Kedves M. 1994. Transmission electron microscopy of the fossil gymnosperm exines. Szeged: Juhász Nyomda.
- Krassilov VA. 1986. New floral structures from the Lower Cretaceous of Lake Baikal area. Review of Palaeobotany and Palynology 47: 9–16. doi:10.1016/0034-6667(86)90003-5.
- Kubitzki K. 1990. Ephedraceae. In: Kubitzki K, ed. The families and genera of vascular plants. I. Pteridophytes and Gymnosperms, 379–382. Berlin: Springer.
- Kurmann MH. 1992. Exine stratification in extant gymnosperms: A review of published transmission electron micrographs. Kew Bulletin 47: 25–39.
- Kurmann MH, Zavada MS. 1994. Pollen morphological diversity in extant and fossil gymnosperms. In: Kurmann MH, Doyle JA, eds. Ultrastructure of fossil spores and pollen, 123–137. London: Royal Botanic Gardens, Kew.
- Land WJG. 1907. Fertilization and embryogeny in Ephedra trifurca. Botanical Gazette 44: 273–292.
- Li Y, Xu Q, Zhao Y, Yang X, Xiao J, Chen H, Lü X. 2005. Pollen indication to source plants in the eastern desert of China. Chinese Science Bulletin 50: 1632–1641.
- Maddison WP, Maddison DR. 2011. Mesquite: A modular system for evolutionary analysis, Version 2.75. <http://mesquiteproject.org>.
- Mehra PN. 1938. The germination of pollen grains in artificial cultures in Ephedra foliata Boiss and Ephedra gerardiana Wall. Proceedings of the Indian Academy of Sciences 8: 218–230.
- Muller J. 1968. Palynology of the Pedawan and Plateau Sandstone formations (Cretaceous–Eocene) in Sarawak, Malaysia. Micropaleontology 14: 1–37.
- Nagy E. 1963. Occurrence of the genus Ephedripites in the Neogene of Hungary. Grana 4: 277–280.
- Narváez PL, Sabino IF. 2008. Palynology of the Las Curtiembres formation (Late Cretaceous, Salta Group Basin), Las Conchas Creek area, northwestern Argentina. Ameghiniana 45: 473–482.
- Niklas KJ. 2015. A biophysical perspective on the pollination biology of Ephedra nevadensis and E. trifurca. The Botanical Review 81: 28–41. doi:10.1007/s12229-014-9148-9.
- Niklas KJ, Buchmann SL, Kerchner V. 1986. Aerodynamics of Ephedra trifurca. I. Pollen grain velocity fields around stems bearing ovules. American Journal of Botany 73: 966–999.
- Niklas KJ, Kerchner V. 1986. Aerodynamics of Ephedra trifurca II. Computer modelling of pollination efficiencies. Journal of Mathematical Biology 73: 966–979.
- Norbäck Ivarsson L. 2014. Pollen morphology in Ephedra (Gnetales) and implications for understanding fossil ephedroid pollen from the Tibetan Plateau, using a phylogenetic approach. Master Thesis, Stockholm University, Stockholm, Sweden.
- Osborn JM. 2000. Pollen morphology and ultrastructure of gymnospermous angiosperms. In: Harley MM, Morton CM,

- Blackmore S, eds. Pollen and spores: Morphology and biology, 163–185. London: Royal Botanical Gardens Kew.
- Osborn JM, Taylor TN, de Lima MR. 1993. The ultrastructure of fossil ephedroid pollen with gnetalean affinities from the Lower Cretaceous of Brazil. Review of Palaeobotany and Palynology 77: 171–184. doi:10.1016/0034-6667(93)90003-D.
- Osborn JM, Taylor TN, Schneider EL. 1991. Pollen morphology and ultrastructure of the Cabombaceae: Correlations with pollination biology. American Journal of Botany 78: 1367–1378.
- Pagel M. 1999a. Inferring the historical patterns of biological evolution. Nature 401: 877–884. doi:10.1038/44766.
- Pagel M. 1999b. The maximum likelihood approach to reconstructing ancestral character states of discrete characters on phylogenies. Systematic Biology 48: 612–622. doi:10.1080/106351599260184.
- Pearson H. 1907. The living *Welwitschia*. Nature 75: 536–537. doi:10.1038/075536a0.
- Pirie MD, Humphreys AM, Antonelli A, Galley C, Linder HP. 2012. Model uncertainty in ancestral area reconstruction: A parsimonious solution? Taxon 61: 652–664.
- R Development Core Team. 2014. R — A language and environment for statistical computing, Version 3.1.1. R Foundation for Statistical Computing, Vienna, Austria, <http://www.r-project.org/>.
- Rowley JR. 1995. Are the endexines of pteridophytes, gymnosperms and angiosperms structurally equivalent? Review of Palaeobotany and Palynology 85: 13–34. doi:10.1016/0034-6667(94)00083-V.
- Rydin C, Bolinder K. 2015. Moonlight pollination in the gymnosperm *Ephedra* (Gnetales). Biology Letters 11: 20140993.
- Rydin C, Friis EM. 2005. Pollen germination in *Welwitschia mirabilis* Hook.f.: Differences between the polyplicate pollen producing genera of the Gnetales. Grana 44: 137–141. doi:10.1080/001713130500230459.
- Rydin C, Khodabandeh A, Endress PK. 2010. The female reproductive unit of *Ephedra* (Gnetales): Comparative morphology and evolutionary perspectives. Botanical Journal of the Linnean Society 163: 387–430. doi:10.1111/boj.2010.163.issue-4.
- Rydin C, Korall P. 2009. Evolutionary relationships in *Ephedra* (Gnetales), with implications for seed plant phylogeny. International Journal of Plant Sciences 170: 1031–1043. doi:10.1086/592013.
- Rydin C, Pedersen KR, Crane PR, Friis EM. 2006a. Former diversity of *Ephedra* (Gnetales): Evidence from early Cretaceous seeds from Portugal and North America. Annals of Botany 98: 123–140. doi:10.1093/aob/mcl078.
- Rydin C, Pedersen KR, Friis EM. 2004. On the evolutionary history of *Ephedra*: Cretaceous fossils and extant molecules. Proceedings of the National Academy of Sciences of the USA 101: 16571–16576. doi:10.1073/pnas.0407588101.
- Rydin C, Wu SQ, Friis EM. 2006b. Liaoxia Cao et S.Q. Wu (Gnetales): Ephedroids from the early Cretaceous Yixian formation in Liaoning, northeastern China. Plant Systematics and Evolution 262: 239–265. doi:10.1007/s00606-006-0481-2.
- Sauquet H, Le Thomas A. 2003. Pollen diversity and evolution in Myristicaceae (Magnoliales). International Journal of Plant Sciences 164: 613–628. doi:10.1086/ijps.2003.164.issue-4.
- Schluter D, Price T, Mooers AØ, Ludwig D. 1997. Likelihood of ancestor states in adaptive radiation. Evolution 51: 1699–1711.
- Scott R. 1960. Pollen of *Ephedra* from the Chinle formation (Upper Triassic) and the genus *Equisetosporites*. Micropaleontology 6: 271–276.
- Shaw CL. 1998. Eocene ephedraceous palynomorphs of Taiwan. Botanical Bulletin of Academia Sinica 39: 60–80.
- Srivastava SK. 1968. Ephedrean pollen from the Upper Cretaceous Edmonton formation of Alberta (Canada) and their paleoecological significance. Canadian Journal of Earth Sciences 5: 211–221. doi:10.1139/e68-022.
- Steeves MW, Barghoorn ES. 1959. The pollen of *Ephedra*. Journal of the Arnold Arboretum 40: 221–255.
- Stover LE. 1964. Cretaceous ephedroid pollen from West Africa. Micropaleontology 10: 145–156.
- Takahashi M. 1995. Ephedroid fossil pollen from the Lower Cretaceous (Upper Albian) of Hokkaido, Japan. Journal of Plant Research 108: 11–15. doi:10.1007/BF02344300.
- Tekleva MV, Krassilov VA. 2009. Comparative pollen morphology and ultrastructure of modern and fossil gnetophytes. Review of Palaeobotany and Palynology 156: 130–138. doi:10.1016/j.revpalbo.2008.12.007.
- Thomas GH, Freckleton RP. 2012. MOTMOT: Models of trait macroevolution on trees. Methods in Ecology and Evolution 3: 145–151. doi:10.1111/j.2041-210X.2011.00132.x.
- Thornhill AH, Popple LW, Carter RJ, Ho SY, Crisp MD. 2012. Are pollen fossils useful for calibrating relaxed molecular clock dating of phylogenies? A comparative study using Myrtaceae. Molecular Phylogenetics and Evolution 63: 15–27. doi:10.1016/j.ympev.2011.12.003.
- Trevisan L. 1980. Ultrastructural notes and considerations on *Ephedrites*, *Eucommioidites* and *Monosulcites* pollen grains from Lower Cretaceous sediments of southern Tuscany (Italy). Pollen et Spores 22: 85–132.
- Van Campo M, Lugardon B. 1973. Structure grenue infratectale de l'ectexine des pollen quelques gymnospermes et angiospermes. Pollen et Spores 15: 171–187.
- Van Emden H. 2008. Statistics for terrified biologists. Oxford: Blackwell.
- Wang X, Zheng SL. 2010. Whole fossil plants of *Ephedra* and their implications on the morphology, ecology and evolution of Ephedraceae (Gnetales). Chinese Science Bulletin 55: 1511–1519. doi:10.1007/s11434-010-3069-8.
- Wang Z. 2004. A new Permian gnetalean cone as fossil evidence for supporting current molecular phylogeny. Annals of Botany 94: 281–288. doi:10.1093/aob/mch138.
- Wetschnig W, Depish B. 1999. Pollination biology of *Welwitschia mirabilis* Hook.f. (Welwitschiaceae, Gnetopsida). Phytion 39: 167–183.
- Wilson LR. 1962. Permian plant microfossils from the Flowerpot formation, Oklahoma. Oklahoma Geological Survey Bulletin 49: 5–50.
- Wodehouse RP. 1933. Tertiary pollen. II. The oil shales of the Eocene Green River formation. Bulletin of the Torrey Botanical Club 60: 479–524.
- Woodhouse RP. 1935. Pollen grains. New York: McGraw-Hill Book Co.
- Yang Y, Geng B-Y, Dilcher DL, Chen Z-D, Lott TA. 2005. Morphology and affinities of an early Cretaceous *Ephedra* (Ephedraceae) from China. American Journal of Botany 92: 231–241. doi:10.3732/ajb.92.2.231.
- Yao Y-F, Xi Y-Z, Geng B-Y, Li C-S. 2004. The exine ultrastructure of pollen grains in Gnetum (Gnetaceae) from China and its bearing on the relationship with the ANITA group. Botanical Journal of the Linnean Society 146: 415–425. doi:10.1111/boj.2004.146.issue-4.
- Zavada MS. 1984. Angiosperm origins and evolution based on dispersed fossil pollen ultrastructure. Annals of the Missouri Botanical Garden 71: 444–463.
- Zhang Y, Xi Y. 1983. Studies on the pollen morphology of Chinese *Ephedra*. Acta Botanica Sinica 25: 418–427.